

VEGETABLE INSECT CONTROL RECOMMENDATIONS (COMMERCIAL)

Crop	Insect	Insecticide	Formulation Per Acre	Lb ai/Acre	Application	Remarks/Precautions	Days to Harvest
ASPARAGUS	Asparagus beetle, Japanese beetle, Grasshoppers	acetamiprid Assail 30 SG	2.5 oz	0.048			1
		carbaryl Sevin XLR Sevin 4F	1 qt 1 qt	1.0	When damage occurs.	DO NOT apply during bloom.	1
		malathion Malathion 57% EC	2 pt	1.25			1
		methomyl (R) Lannate 2.4 LV Lannate 90 SP	1.5-3 pt 1/2-1 lb	0.45-0.9			1
		permethrin (R) Pounce 25 WP	3.2-6.4 oz	0.05-0.1			1
		spinetoram Radiant 1 SC	4-8 oz	0.03-0.063		After harvest of ferns only.	3
		spinosad SpinTor 2 SC	4-6 oz	0.062-0.094		For protection of ferns only.	60
		<i>Bacillus thuringiensis</i> Biobit, Dipel, Javelin, etc.	See label for rates.				0
	Beet armyworm, Cutworms, Yellow-striped armyworm	chlorantraniliprole Coragen 1.67 SC Shenzi 400SC	3.5-7.5 fl oz 1.7-3.8 fl oz	0.045-0.098		DO NOT use an adjuvant with Coragen.	1
		Cyantraniliprole Exirel 0.83 EC	7-13.5 fl oz	0.45-0.88			1
		methomyl (R) Lannate 2.4 LV Lannate 90 SP	1 1/2-3 pt 1/2-1 lb	0.45-0.9	When damage occurs.		1
BEANS (snap or lima)	Bean leaf beetle, Mexican bean beetle	acetamiprid Assail 30 SG	2.5-5.3 oz	0.048-0.10			7
		bifenthrin (R) Brigade 2 EC	2.6-6.4 fl oz	0.04-0.1			3
		carbaryl Sevin XLR Sevin 4F	1/2-1 qt 1/2-1 qt	0.5-1	When damage occurs.	DO NOT apply during bloom.	3
		esfenvalerate (R) Asana XL 0.66 EC	2.9-5.8 oz	0.015-0.03		Asana is not registered on lima beans.	3
		gamma-cyhalothrin Proaxis 0.5EC	2.56-3.84 oz	0.01-0.015			7
		lambda-cyhalothrin (R) Karate Z	1.28-1.92 oz	0.02-0.03			7
		methomyl (R) Lannate 90 SP	0.5 lb	0.45			1
		Novaluron Rimon 0.83 EC	9-12 oz	0.058-0.077		Controls immature mexican bean beetles only.	1
		zeta-cypermethrin (R) Mustang Maxx 0.8 EC	3.2-4.0 oz	0.02-0.025			1

VEGETABLE INSECT CONTROL RECOMMENDATIONS (COMMERCIAL)

Crop	Insect	Insecticide	Formulation Per Acre	Lb ai/Acre	Application	Remarks/Precautions	Days to Harvest
BEANS (snap or lima) (cont.)	Cabbage looper	<i>Bacillus thuringiensis</i> Agree WG		2		Loopers rarely require management on Arkansas snap bean.	
		chlorantraniliprole Shenzi 400SC Coragen	1.7-3.8 fl oz 3.5-7.5 fl oz	0.045-0.098			1
		esfenvalerate (R) Asana XL 0.66 EC	5.8-9.6 oz	0.03-0.05		Asana is not registered on lima beans.	3
		methomyl (R) Lannate 90 SP Lannate 2.4 LV	0.5-1 lb 1 1/2-3 pt	0.45-0.9			3
		methoxyfenozide Intrepid 2F	8-16 oz	0.16-0.25			7
Corn earworm	bifenthrin (R) Brigade 2 EC		2.6-6.4 fl oz	0.04-0.1		In spring planted snap bean, scouting (300 sweeps per 50 acres) can be used to reduce insecticide applications.	3
		carbaryl Sevin 5% D	40 lb	2.0	When first pods are 1 inch long and weekly thereafter.	Fall-planted snap beans will generally require corn earworm management. DO NOT apply during bloom.	3
		chlorantraniliprole Coragen 1.67 SC Shenzi 400SC	3.5-7.5 fl oz 1.7-3.8 fl oz	0.045-0.098			1
		Cyantraniliprole Exirel 0.83 EC	7-13.5 fl oz	0.45-0.88			1
		esfenvalerate (R) Asana XL 0.66 EC	5.8-9.6 oz	0.03-0.05		DO NOT feed treated vines. Asana is not registered on lima beans.	3
		gamma-cyhalothrin Proaxis 0.5EC	2.56-3.84 oz	0.01-0.015			7
		lambda-cyhalothrin (R) Karate Z	1.28-1.92 oz	0.02-0.03			7
		methomyl (R) Lannate 90 SP Lannate 2.4 LV	0.5 lb 1 1/2 pt	0.45			1
		methoxyfenozide Intrepid 2F	10-16 oz	0.16-0.25			7
		spinetoram Radiant 1 SC	4-8 oz	0.03-0.063			3 (succulent); 28 (dried)
		spinosad Blackhawk	1.7-3.3 oz	0.04-0.07			3 (succulent); 28 (dried)
		zeta-cypermethrin (R) Mustang Maxx 0.8 EC	3.2-4.0 oz	0.02-0.025			1
		zeta-cypermethrin +bifenthrin (R) Hero	10.3 oz	0.1			3

VEGETABLE INSECT CONTROL RECOMMENDATIONS (COMMERCIAL)

Crop	Insect	Insecticide	Formulation Per Acre	Lb ai/Acre	Application	Remarks/Precautions	Days to Harvest
BEANS (snap or lima) (cont.)	European corn borer	chlorantraniliprole Coragen 1.67 SC Shenzi 400SC	3.5-7.5 fl oz 1.7-3.8 fl oz	0.045-0.098		Corn borers rarely require management on Arkansas snap bean.	1
		Cyantraniliprole Exirel 0.83 EC	7-13.5 fl oz	0.45-0.88			1
		zeta-cypermethrin + bifenthrin (R) Hero	10.3 oz	0.1			3
Mites	abamectin Agri-Mek SC		1.75-3.5 fl oz	0.01-0.05	For control of spider mites.	DO NOT make more than 2 applications. Must be mixed with an adjuvant.	7
	acequinocyl Kanemite 15 SC		31 fl oz	0.3		DO NOT make more than 2 applications.	7
	fenpyroximate Portal 0.4 EC		2 pt	0.1		Snap bean only. DO NOT make more than 2.	1 (Snap bean)
	bifenthrin (R) Brigade 2 EC		6.4 fl oz	0.1			3
	dimethoate Dimethoate 4EC		1 pt	0.5			0
Seedcorn maggot, Wireworms	bifenthrin Empower 1.15 G		3.5-8.7 lb			Apply preplant broadcast in the top 1 to 3 inches of soil.	9
	phorate (R) Thimet 20G			7.6	Distribute 4.5-7.0 oz of the granules evenly in 1,000 row feet over the top of the row at planting time (30-inch row minimum).	DO NOT feed foliage of treated beans. DO NOT place Thimet in contact with seed.	
Stink bugs	bifenthrin (R) Brigade 2 EC		2.6-6.4 fl oz	0.04-0.1			3
	lambda-cyhalothrin (R) Karate Z		1.28-1.92 oz	0.02-0.03			7
	zeta-cypermethrin (R) Mustang Maxx 0.8 EC		3.2-4.0 oz	0.02-0.025			1
	zeta-cypermethrin + bifenthrin (R) Hero		10.3 oz	0.1			3
BEETS	Flea beetles, webworms	carbaryl Sevin 4F Sevin XLR Plus	1 qt 1 qt	1.0	When insects appear in damaging numbers.	DO NOT apply during bloom.	7
		imidacloprid Prey 1.6 F	3.5 oz	0.044			7
		zeta-cypermethrin (R) Mustang Maxx 0.8 EC	1.8-4.0 oz	0.011-0.025			1

VEGETABLE INSECT CONTROL RECOMMENDATIONS (COMMERCIAL)

Crop	Insect	Insecticide	Formulation Per Acre	Lb ai/Acre	Application	Remarks/Precautions	Days to Harvest
CARROTS	Flea beetles, celery worm, leafhoppers	beta-cyfluthrin + imidacloprid Leverage 360	2.8 oz	0.066			7
		carbaryl Sevin 4F Sevin XLR Plus	1 qt 1 qt	1.0	When insects appear in damaging numbers.	Avoid application when foliage is wet. DO NOT apply during bloom.	7
		imidacloprid Prey 1.6 F	3.5 oz	0.044			7
		malathion Malathion 57% EC	2 pt	1.25	Use only for leafhoppers.		7
		sulfoxaflor Transform WG	1.5-2.75	0.047-0.086		Leafhopper control only. Check label for crop rotation restrictions.	7
		zeta-cypermethrin (R) Mustang Maxx 0.8 EC	1.8-4.0 oz	0.011-0.025			1
COLE CROPS (Broccoli, Brussels sprouts, cabbage and cauliflower)	Aphids	acetamiprid Assail 30 SG	2.5 oz	0.048			1
		acetamiprid + bifenthrin Savoy	4.9-12.9 oz			Maximum of 19 oz a season.	7
		beta-cyfluthrin + imidacloprid Leverage 360	3 oz	0.07			7
		bifenthrin (R) Brigade 2 EC	2.6-6.4 fl oz	0.04-0.1			7
		flonicamid Beleaf 50SG	2-2.8 oz	0.062-0.089			0
		flupyradifurone Sivanto Prime	7-14 fl oz	0.09-0.18		Except green peach aphid. Do not apply more than 28 fl oz in a season.	1
		imidacloprid Admire Pro 4.6 F Prey 1.6 F	3.75 oz 1.3 fl oz	0.047			7
		lambda-cyhalothrin (R) Karate Z	1.28-1.92 fl oz	0.02-0.03			1
		zeta-cypermethrin (R) Mustang Maxx 0.8 EC	2.24-4.0 oz	0.014-0.025			1
		zeta-cypermethrin + bifenthrin Hero	10.3 oz	0.1			3
Caterpillars (cabbage looper, diamondback, imported and cross-striped cabbage worm)	acetamiprid + bifenthrin Savoy	9.6-12.9 oz				Use on diamond back moth only. Maximum of 19 oz a season.	7
		Bacillus thuringiensis (Dipel, Javelin, Agree, Crymax)			Check label.	Slow acting but works well.	0

VEGETABLE INSECT CONTROL RECOMMENDATIONS (COMMERCIAL)

Crop	Insect	Insecticide	Formulation Per Acre	Lb ai/Acre	Application	Remarks/Precautions	Days to Harvest	
COLE CROPS (Broccoli, Brussels sprouts, cabbage and cauliflower) (cont.)	Caterpillars (cabbage looper, diamondback, imported and cross-striped cabbage worm) (cont.)	beta-cyfluthrin + imidacloprid Leverage 360 bifenthrin (R) Brigade 2 EC chlorantraniliprole Coragen 1.67 SC Shenzi 400SC chlorantraniliprole + lambda-cyhalothrin (R) Voliam Xpress Cyantraniliprole Exirel 0.83 EC esfenvalerate (R) Asana XL 0.66 EC lambda-cyhalothrin (R) Karate Z methomyl (R) Lannate 2.4 LV Lannate SP methoxyfenozide Intrepid 2F permethrin (R) Ambush 25 W Ambush 2.0 EC Pounce 25 WP spinosad SpinTor 2 SC zeta-cypermethrin (R) Mustang Maxx 0.8 EC zeta-cypermethrin + bifenthrin Hero	3 oz 2.6-6.4 fl oz 3.5-7.5 fl oz 1.7-3.8 fl oz 6-9 oz 7-13.5 fl oz 5.8-9.6 oz 1.28-1.92 fl oz 1/2-3 pt 1/2-1 lb 4-8 oz 3.2-6.4 oz 3.2-12.8 oz 3.2-6.4 oz 1.5-10 fl oz 2.24-4.0 oz 10.3 oz	0.07 0.04-0.1 0.045-0.098 0.06-0.09 0.45-0.88 0.03-0.05 0.02-0.03 0.45-0.9 0.06-0.12 0.05-0.1 0.023-0.156 0.014-0.025 0.1			Follow label directions. DO NOT feed treated plants to livestock.	7 7 3 3 1 3 1 1 1 1 1 1 1 7
	Cutworms	beta-cyfluthrin Baythroid XL bifenthrin (R) Brigade 2 EC cypermethrin Holster 2.5 EC diazinon (R) Diazinon AG500 Diazinon AG600 WBC esfenvalerate (R) Asana XL 0.66 EC zeta-cypermethrin (R) Mustang Maxx 0.8 EC	0.8-1.6 oz 2.1-6.4 2.5-5 oz 2-4 qt 51-102 fl oz 5.8-9.6 oz 3.2-4.0 oz	0.007-0.013 0.033-0.01 0.05-0.01 2.0-4.0 Apply when cabbage loopers are small. Broadcast just before planting and incorporate.		Follow label directions. Follow label directions.	0 7 1 3 1	

VEGETABLE INSECT CONTROL RECOMMENDATIONS (COMMERCIAL)

Crop	Insect	Insecticide	Formulation Per Acre	Lb ai/Acre	Application	Remarks/Precautions	Days to Harvest	
COLE CROPS (Broccoli, Brussels sprouts, cabbage and cauliflower) (cont.)	Harlequin bug, stink bugs	acetamiprid + bifenthrin Savoy	9.6-12.9 oz			Maximum of 19 oz a season.	7	
		bifenthrin (R) Brigade 2 EC	2.1-6.4 oz	0.033-0.10			7	
		beta-cyfluthrin + imidacloprid Leverage 360	3 oz	0.07			7	
		carbaryl Sevin XLR Plus	1-2 qt	1.2		Follow label directions. DO NOT apply during bloom.	14	
		Sevin 4F	1-2 qt					
		clothianidin Belay 50 WDG	3 - 4 oz	0.05-0.067			7	
		dinotefuran Venom 70 SG	3-4 oz	0.05-0.18			1	
CUCURBITS (Cantaloupe, cucumber, melons, pumpkin, squash, summer squash and watermelon)	Aphids	acetamiprid Assail 30 SG	2.5 oz-4 oz	0.048-0.077			0	
		acetamiprid + bifenthrin Savoy	9.6-12.9 oz			Maximum of 38.7 oz a season.	3	
		flonicamid Beleaf 50SG	2-2.8 oz	0.062-0.089			0	
		flupyradifurone Sivanto Prime	7-14 fl oz	0.09-0.18		DO NOT use foliar applications in muskmelon, cantaloupe, or honeydew melon.	1	
		sulfoxaflor Transform WG	0.75	0.023		Check label for crop rotation restrictions.	1	
	Cucumber beetles	acetamiprid Assail 30 SG	2.5 oz-5.3 oz	0.05-0.1			1	
		acetamiprid + bifenthrin Savoy	6-12.9 oz			Maximum of 38.7 oz a season.	3	
		beta-cyfluthrin Baythroid XL	2.4-2.8 oz	0.019-0.022			0	
		bifenthrin (R) Brigade 2 EC	2.6-6.4 fl oz	0.04-0.1			3	
		carbaryl Sevin XLR Plus	1 qt	1.0	When insects first appear.	Sevin is highly toxic to honeybees. Avoid excessive applications. DO NOT apply when foliage is wet. DO NOT apply during bloom.	3	
		Sevin 4F	1 qt					
		clothianidin Belay 50 WDG	3 – 4 oz	0.05-0.067		DO NOT use adjuvant with foliar applications. DO NOT use after 4th true leaf.	-	
		dinotefuran Venom 70 SG	3-4 oz	0.05-0.18			1	
		Scorpion 35 SL	2-7 fl oz					

VEGETABLE INSECT CONTROL RECOMMENDATIONS (COMMERCIAL)

Crop	Insect	Insecticide	Formulation Per Acre	Lb ai/Acre	Application	Remarks/Precautions	Days to Harvest
CUCURBITS (Cantaloupe, cucumber, melons, pumpkin, squash, summer squash and watermelon)	Cucumber beetles (cont.)	imidacloprid Admire Pro 4.6 F	7-10.5 fl oz	0.28-0.38		Admire applied into soil is highly effective against cucumber beetles and can reduce seedling loss. Soil Application Only.	21
		lambda-cyhalothrin (R) Karate Z	1.28-1.92 fl oz	0.02-0.03			1
		malathion Malathion 5 EC	1.5-2.75 pt	0.94-1.7		Use no more than 1.5 pints on winter squash or pumpkins or 2 pints on watermelons.	1
		methomyl (R) Lannate 2.4 LV Lannate SP	1 1/2-3 pt 1/2-1 lb	0.45-0.9		DO NOT use Lannate on pumpkins.	1 (lower rate) 3 (higher rate)
		zeta-cypermethrin (R) Mustang Maxx 0.8 EC	2.8-4.0 oz	0.0175-0.025			1
		zeta-cypermethrin + bifenthrin Hero	10.3 oz	0.1			3
Flea beetles Tobacco budworm Beet armyworm		methomyl (R) Lannate 2.4 LV Lannate SP	1 1/2-3 pt 1/2-1 lb	0.45-0.9	When insects first appear.	One day up to 2 pt, over 2 pt 3 days. DO NOT apply to pumpkins.	1-3
		methoxyfenozide Intrepid 2F	4-10 oz	0.06-0.16		Not for control of tobacco budworm.	3
		spinosad SpinTor 2 SC	4-8 fl oz	0.0625-0.125		Not labeled for flea beetles.	3
Mites (Mites are primarily a problem on cucumber and watermelon)		abamectin Agri-Mek 0.7 SC	1.75-3.5 oz	0.01-0.019		Must be mixed with an adjuvant (NIS surfactant).	7
		bifenazate Acramite 50 WS	0.75-1 lb	0.375-0.5		Only one application per season.	3
		extazole Zeal 2.88 SC	4-6 fl oz	0.09-0.135		Does not kill Adults. DO NOT make more than 1 application per season.	7
		Fenpyroximate Portal	2 pt				1
		spiromesifen Oberon 2 SC	7-8.5 oz	0.11-0.13			7
Pickleworm, Melonworm, Cutworms		acetamiprid + bifenthrin Savoy	6-12.9 oz			Maximum of 38.7 oz a season.	3
		beta-cyfluthrin Baythroid XL	1.6-2.4 oz	0.013-0.019			0
		bifenthrin (R) Brigade 2 EC	2.6-6.4 fl oz	0.04-0.1			3
		carbaryl Sevin XLR Plus Sevin 4F	1 qt 1 qt	1.0	When insects or damage first appears.	Avoid excessive applications. DO NOT apply when foliage is wet. DO NOT apply during bloom.	3
		chlorantraniliprole Coragen 1.67 SC Shenzi 400SC	2.0-7.5 fl oz 1.0-3.8 fl oz	0.026-0.098			3
		Cyantraniliprole Exirel 0.83 EC	7-13.5 fl oz	0.45-0.88			1
							1

VEGETABLE INSECT CONTROL RECOMMENDATIONS (COMMERCIAL)

Crop	Insect	Insecticide	Formulation Per Acre	Lb ai/Acre	Application	Remarks/Precautions	Days to Harvest
CUCURBITS (Cantaloupe, cucumber, melons, pumpkin, squash, summer squash and watermelon) (cont.)	Pickleworm, Melonworm, Cutworms (cont.)	esfenvalerate (R) Asana XL 0.66 EC	5.8-9.6 oz	0.03-0.05		Asana may cause some phytotoxicity to zucchini squash.	3
		lambda-cyhalothrin (R) Karate Z	1.28-1.92 oz	0.02-0.03			1
		methoxyfenozide Intrepid 2F	4-10 fl oz	0.06-0.15			3
		permethrin (R) Ambush 2.0 EC Pounce 3.2 EC	6.4-12.8 oz 4-8 oz	0.1-0.2	When insects or damage first appear.		0
		spinetoram Radiant 1 SC	5-10 oz	0.038-0.079			3
		spinosad SpinTor 2 SC	4-8 oz	0.062-0.125		Follow label directions. SpinTor does not control squash bug.	3
		zeta-cypermethrin + bifenthrin Hero	10.3 oz	0.1			3
	Squash Bug (Rarely a pest of cucumber or watermelon)	acetamiprid + bifenthrin Savoy	12.9 oz			Maximum of 38.7 oz a season.	3
		beta-cyfluthrin Baythroid XL	1.6-2.4 oz	0.013-0.019			0
Squash Vine Borer (Only attacks squash and pumpkin)	bifenthrin (R) Brigade 2 EC	2.6-6.4 fl oz	0.04-0.1			Bifenthrin is the only registered insecticide that provides moderate but effective control of squash bug nymphs. Almost no control should be expected of adult squash bugs.	3
	carbaryl Sevin XLR Plus Sevin 4F	1 qt 1 qt	1.0	When insects or damage first appear.		Avoid excessive applications. DO NOT apply when foliage is wet. DO NOT apply during bloom.	3
	esfenvalerate (R) Asana XL 0.66 EC	5.8-9.6 oz	0.03-0.05			Asana may cause some phytotoxicity to zucchini squash.	3
	lambda-cyhalothrin (R) Karate Z	1.28-1.92 fl oz	0.02-0.03				1
	permethrin (R) Ambush 2.0 Pounce 3.2	6.4-12.8 oz 4-8 oz	0.1-0.2	When insects or damage first appear.			0
	zeta-cypermethrin + bifenthrin Hero	10.3 oz					3
	acetamiprid Assail 30 SG	5 oz	0.1			Not generally a serious issue in large plantings of squash or pumpkin.	0
	acetamiprid + bifenthrin Savoy	6-12.9 oz	0.1			Consider a blue hubbard squash trap crop for control.	3
	bifenthrin (R) Brigade 2 EC	2.6-6.4 fl oz	0.04-0.1				3
	chlorantraniliprole Coragen 1.67 SC	3.5-7.5 fl oz	0.045-0.098				3
	lambda-cyhalothrin (R) Karate Z	1.28-1.92 fl oz	0.02-0.03				1

VEGETABLE INSECT CONTROL RECOMMENDATIONS (COMMERCIAL)

Crop	Insect	Insecticide	Formulation Per Acre	Lb ai/Acre	Application	Remarks/Precautions	Days to Harvest
CUCURBITS (Cantaloupe, cucumber, melons, pumpkin, squash, summer squash and watermelon) (cont.)	Whiteflies	acetamiprid Assail 30 SG	2.5 oz-4 oz	0.048-0.077		Insecticide use will not effectively reduce high populations of whiteflies. Removal of alternate host plants (Weeds) along with field plowing immediately after harvest aids in population suppression.	0
		acetamiprid + bifenthrin Savoy	6-12.9 oz				3
		chlorantraniliprole Coragen 1.67 SC Shenzi 400SC	5-7.5 fl oz 2.5-3.5 fl oz	0.065-0.098	Use on nymphs only. See label for remarks.		1
		Cyantraniliprole Exirel 0.83 EC	13.5-20 fl oz	0.088-0.133			1
		pyriproxyfen Knack 0.86 EC	8-10 oz	0.05-0.067			7
		sulfoxaflor Transform WG	2.0-2.25	0.063-0.071			1
		thiomethoxam Actara 25 WDG	3-5.5 oz	0.047-0.086	Follow label directions.		0
EGGPLANT	Aphids	acetamiprid Assail 30 SG	2.5 oz-4 oz	0.048-0.077			0
		acetamiprid + bifenthrin Savoy	4.9-9.6 oz			Follow label directions.	7
		beta-cyfluthrin + imidacloprid Leverage 360	3.8-4.1 oz	0.89-0.096			7
		flonicamid Beleaf 50 SG	2.8-4.28 oz	0.089-0.133			0
		flupyradifurone Sivanto Prime	7-12 fl oz	0.09-0.15			1
		imidacloprid Admire Pro 4.6 F	1.3-2.2 fl oz	0.05-0.08			0
		sulfoxaflor Transform WG	0.75-1.0	0.023-0.031		Check label for crop rotation restrictions.	1
		thiamethoxam Actara 25 WDG	2-3 oz	0.03-0.047			0
Caterpillars		beta-cyfluthrin Baythroid XL	1.6-2.8 oz	0.013-0.022		Use higher rates for loopers, pinworms and armyworms.	7
		beta-cyfluthrin + imidacloprid Leverage 360	3.8-4.1 oz	0.89-0.096			7
		chlorantraniliprole Coragen 1.67 SC Shenzi 400SC	3.5-7.5 fl oz 1.0-3.8 fl oz	0.045-0.098			1
		Cyantraniliprole Exirel 0.83 EC	7-13.5 fl oz	0.45-0.88			1
		chlorantraniliprole + lambda-cyhalothrin (R) Voliam Xpress	6-9 oz	0.06-0.09			5
		methoxyfenozide Intrepid 2F	4-16 fl oz	0.06-0.25			1

VEGETABLE INSECT CONTROL RECOMMENDATIONS (COMMERCIAL)

Crop	Insect	Insecticide	Formulation Per Acre	Lb ai/Acre	Application	Remarks/Precautions	Days to Harvest
EGGPLANT (cont.)	Flea beetles, Colorado potato beetle	beta-cyfluthrin Baythroid XL	2.8 oz	0.022			7
		beta-cyfluthrin + imidacloprid Leverage 360	3.8-4.1 oz	0.89-0.096			7
		bifenthrin (R) Brigade 2 EC	2.1-6.4 fl oz	0.033-0.1			7
		carbaryl Sevin XLR Plus Sevin 4F	1/2-2 qt 1/2-2 qt	0.5-2.0		DO NOT apply during bloom. Use higher rates for Colorado potato beetle.	3
		esfenvalerate (R) Asana XL 0.66 EC	5.8-9.6 oz	0.03-0.05			7
		imidacloprid Admire Pro 4.6 F	1.3-2.2 fl oz	0.05-0.08			0
		zeta-cypermethrin (R) Mustang Maxx 0.8 EC	2.24-4.0 oz	0.014-0.025			1
Mites		abamectin Agri-Mek 0.7 SC	1.75-3.5 fl oz	0.01-0.019		Must be mixed with an adjuvant.	7
		fenbutatin-oxide (R) Vendex 50 WP	2-3 lb	1.0-1.5			3
		Fenpyroximate Portal	2 pt			Apply in a minimum of 20 gallons per acre	1
		spiromesifen Oberon 2 SC	7-8.5 oz	0.11-0.13			1
Leafminers		permethrin (R) Ambush 2.0 EC Pounce 25 WP	6-12 oz 6.4-9.6 oz	0.1-0.2			3
		spinosad SpinTor 2 SC	6-8 oz	0.094-0.125		Follow label directions.	1
GREENS (collards)	Caterpillars	<i>Bacillus thuringiensis</i>					
		beta-cyfluthrin Baythroid XL	1.6-3.2 oz	0.013-0.025		Use higher rates for earworms, armyworms and European corn borers.	0
		beta-cyfluthrin + imidacloprid Leverage 360	3 oz	0.07			7
		chlorantraniliprole Coragen 1.67 SC Shenzi 400SC	3.5-7.5 fl oz 1.7-3.8 fl oz	0.045-0.098			3
		Cyantraniliprole Exirel 0.83 EC	7-13.5 fl oz	0.45-0.88			1
		esfenvalerate (R) Asana XL 0.66EC	5.8-9.6 oz	0.03-0.05			7
		indoxacarb Avaunt 30 WDG	2.5-3.5 oz	0.045-0.065			3

VEGETABLE INSECT CONTROL RECOMMENDATIONS (COMMERCIAL)

Crop	Insect	Insecticide	Formulation Per Acre	Lb ai/Acre	Application	Remarks/Precautions	Days to Harvest
GREENS (collards)	Caterpillars (cont.)	methoxyfenozide Intrepid 2F	4-16 oz	0.06-0.25			1
		permethrin (R) Ambush 2.0 EC	3.2-6.4 oz	0.05-0.1			1
		Pounce 25 WP	3.2-9.6 oz				
GREENS (kale and mustard)	Caterpillars	zeta-cypermethrin (R) Mustang Maxx 0.8 EC	2.24-4.0 oz	0.014-0.025			1
		<i>Bacillus thuringiensis</i>					
		beta-cyfluthrin Baythroid XL	16-3.2 oz	0.013-0.025		Use higher rates for earworms and armyworms.	0
		beta-cyfluthrin + imidacloprid Leverage 360	3 oz	0.07			7
		chlorantraniliprole Coragen 1.67 SC	3.5-7.5 fl oz	0.045-0.098			3
		Shenzi 400SC	1.7-3.8 fl oz				
		Cyantraniliprole Exirel 0.83 EC	7-13.5 fl oz	0.45-0.88			1
		cypermethrin Holster 2.5 EC	2.5-5.0 oz	0.05-0.1			1
		methoxyfenozide Intrepid 2F	4-8 fl oz	0.06-0.125			1
GREENS (turnips)	Caterpillars	spinosad SpinTor 2 SC	3-6 oz	0.047-0.094			1
		zeta-cypermethrin (R) Mustang Maxx 0.8 EC	2.24-4.0 oz	0.014-0.025			1
		<i>Bacillus thuringiensis</i>					
		beta-cyfluthrin Baythroid XL	1.6-3.2 oz	0.013-0.025		Use higher rates for corn earworms and armyworms.	0
		beta-cyfluthrin + imidacloprid Leverage 360	3 oz	0.07			7
		chlorantraniliprole Coragen	3.5-7.5 fl oz	0.045-0.098			1
		Shenzi 400SC	1.7-3.8 fl oz				
		methoxyfenozide Intrepid 2F	8-16 oz	0.12-0.25			1
		permethrin (R) Pounce 3.2 EC	2-4 oz	0.05-0.1			1

VEGETABLE INSECT CONTROL RECOMMENDATIONS (COMMERCIAL)

Crop	Insect	Insecticide	Formulation Per Acre	Lb ai/Acre	Application	Remarks/Precautions	Days to Harvest
HOPS	Aphids and Leafhoppers	imidacloprid Admire Pro 4.6 F	2.8 fl oz	0.1			28
		pymetrozine Fulfill 50 WDG	4-6 oz	0.125-0.187		Aphids only, will not control leaf hoppers.	14
		spirotetramat Movento	5-6 fl oz	0.08-0.09		Do not exceed 12.5 fl oz per season.	7
		Malathion Malathion 5 EC Malathion 8 EC	1 pt 0.63 pt	1.25			10
		Pyrethrins Pyganic EC 1.4 II Pyganic EC 5 II	16-64 fl oz 4.5-17 fl oz	0.175-0.7		OMRI approved. Do not expect residual control.	0
	Japanese Beetle	bifenthrin (R) Brigade 2 EC	3.8-6.4 fl oz	0.06-0.1			14
		imidacloprid Admire Pro 4.6 F	2.8 fl oz	0.1			28
	Armyworms, Cutworms, Loopers, Leafrollers, and other caterpillars	Bacillus thuringiensis Bt Containing products					
		bifenthrin (R) Brigade 2 EC	3.8-6.4 fl oz	0.06-0.1			14
		chlorantraniliprole Coragen 1.67 SC Shenzi 400SC	3.5-7.5 fl oz 1.7-3.8 fl oz	0.045-0.098			0
		spinosad Entrust 2 SC	4-6 oz	0.063-0.094		OMRI approved.	1
		spinetoram Delegate 25 WG	2.5-4 oz				3
Spider Mites	Spider Mites	abamectin Agri-Mek 0.7 SC	1.75-3.5 fl oz	0.01-0.019		Must be mixed with an adjuvant (NIS surfactant).	7
		acequinocyl Kanemite 15 SC	31 fl oz	0.3			7
		bifenazate Acramite 50 WS	0.75-1 lb	0.375-0.5		Only one application per season.	14
		extazole Zeal 72 WSP	3-4 oz	0.135-0.18		Does not kill Adults. DO NOT make more than 1 application per season.	7
		fenpyroximate Portal 0.4 EC	2 pt	0.1			3
OKRA	Aphids	acetamiprid + bifenthrin Savoy	4.9-9.6 oz			Follow label directions.	7
		flonicamid Beleaf 50 SC	2.8-4.28 oz	0.089-0.133			0
		imidacloprid Admire Pro 4.6 F	1.3-2.2 fl oz	0.05-0.08			0
		sulfoxaflor Transform WG	0.75-1.0	0.023-0.031		Check label for crop rotation restrictions.	1

VEGETABLE INSECT CONTROL RECOMMENDATIONS (COMMERCIAL)

Crop	Insect	Insecticide	Formulation Per Acre	Lb ai/Acre	Application	Remarks/Precautions	Days to Harvest
OKRA (cont.)	Corn earworm	bifenthrin (R) Brigade 2 EC	2.1-6.4 oz	0.033-0.1			7
		chlorantraniliprole Coragen 1.67 SC	3.5-7.5 fl oz	0.045-0.098			1
		Shenzi 400SC	1.7-3.8 fl oz				
		Cyantraniliprole Exirel 0.83 EC	7-13.5 fl oz	0.45-0.88			1
		methoxyfenozide Intrepid 2F	4-16 oz	0.06-0.25			1
		spinosad SpinTor 2 SC	3-6 oz	0.047-0.094			1
ONION	Thrips	acetamiprid Assail 30 SG	2.1 oz-3.4 oz	0.040-0.065			7
		lambda-cyhalothrin (R) Karate Z	1.28-1.92 fl oz	0.02-0.03			14
		malathion Malathion 57% EC	1 1/2 pt	0.9		Highly toxic to honeybees.	3
		spinetoram Radiant 1 SC	6-8 oz	0.047-0.063			3
		sulfoxaflor Transform WG	2.75	0.086		Check label for crop rotation restrictions.	7
		zeta-cypermethrin (R) Mustang Maxx 0.8 EC	2.24-4.0 oz	0.014-0.025			7
PEAS (Garden or English)	Aphids	acetamiprid + bifenthrin Savoy	4.9-9.6 oz			Follow label directions.	7
		esfenvalerate (R) Asana XL 0.66 EC	2.9-5.8 oz	0.015-0.05			3
		imidacloprid Admire Pro 4.6F	1.2 fl oz	0.05			7
		lambda-cyhalothrin (R) Karate Z	1.28-1.92 fl oz	0.02-0.03			7
		malathion Malathion 57% EC 5 lb/gal	1 1/2 pt	0.9	When insects appear.	Seven days if forage is to be fed to livestock.	3
PEPPERS	Aphids	acetamiprid + bifenthrin Savoy	4.9-9.6 oz			Follow label directions.	7
		beta-cyfluthrin + imidacloprid Leverage 360	3.8-4.1 oz	0.89-0.096			7
		flonicamid Beleaf 50 SG	2.0-2.8 oz	0.026-0.69			0
		imidacloprid Admire Pro 4.6F	1.2 fl oz	0.047			0
		methomyl (R) Lannate 2.4 LV Lannate SP	1 1/2 pt 1/2 lb	0.45			3

VEGETABLE INSECT CONTROL RECOMMENDATIONS (COMMERCIAL)

Crop	Insect	Insecticide	Formulation Per Acre	Lb ai/Acre	Application	Remarks/Precautions	Days to Harvest
PEPPERS (cont.)	Aphids (cont.)	sulfoxaflor Transform WG	0.75-1.0	0.023-0.031		Check label for crop rotation restrictions.	1
		thiamethoxam Actara 25 WDG	2-3 oz	0.033-0.047			0
	Cutworms	bifenthrin (R) Brigade 2 EC	2.1-6.4 oz	0.033-0.1			7
		lambda-cyhalothrin (R) Karate Z	0.96-1.6 fl oz	0.015-0.025			5
		zeta-cypermethrin (R) Mustang Maxx 0.8 EC	1.28-4.0 oz	0.008-0.025			1
		zeta-cypermethrin + bifenthrin Hero	10.3 oz	0.1			7
	Flea beetles, Colorado potato beetle, earworm, European corn borer, hornworms	acetamiprid + bifenthrin Savoy	3.6-6.0 oz	0.013-0.022		Follow label directions. Colorado potato beetle only.	7
		beta-cyfluthrin Baythroid XL	1.6-2.8 oz				7
		beta-cyfluthrin + imidacloprid Leverage 360	3.8-4.1 oz	0.89-0.096			7
		bifenthrin (R) Brigade 2 EC	2.1-6.4 oz	0.033-0.1			7
CORN (cont.)	carbaryl Sevin 4F Sevin XLR Plus		1-2 qt 1-2 qt	1.0-2.0	When insects appear.	DO NOT apply during bloom.	3
		chlorantraniliprole Coragen 1.67 SC Shenzi 400SC	2.0-7.5 fl oz 1.0-3.8 fl oz	0.026-0.098			1
		Cyantraniliprole Exirel 0.83 EC	7-13.5 fl oz	0.45-0.88			1
		esfenvalerate (R) Asana XL 0.66 EC	5.8-9.6	0.03-0.05			7
		lambda-cyhalothrin (R) Karate Z	1.28-1.92 fl oz	0.02-0.03			5
		methoxyfenozide Intrepid 2F	4-16 oz	0.08-0.25		European corn borer and hornworm control.	1
		permethrin (R) Pounce 25 WP	6.4-12.8 oz	0.1-0.2			3
		spinetoram Radiant 1 SC	5-10 fl oz	0.04-0.078			1
		spinosad SpinTor 2 SC	3-8 fl oz	0.047-0.125			1
		zeta-cypermethrin (R) Mustang Maxx 0.8 EC	2.24-4.0 oz	0.014-0.025			1
CUCUMBERS (cont.)	zeta-cypermethrin + bifenthrin Hero		4.0-10.3 oz	0.04-0.1			7

VEGETABLE INSECT CONTROL RECOMMENDATIONS (COMMERCIAL)

Crop	Insect	Insecticide	Formulation Per Acre	Lb ai/Acre	Application	Remarks/Precautions	Days to Harvest
PEPPERS (cont.)	Mites	abamectin (R) Agri-Mek 0.7 SC	1.75-3.5 fl oz	0.0094- 0.01875		Use only two times per season. Must be mixed with an adjuvant (NIS surfactant).	7
		spiromesifen Oberon 2 SC	7-8.5 oz	0.11-0.13			1
	Other caterpillars	bifenthrin (R) Brigade 2 EC	2.1-6.4 oz	0.033-0.1			7
		beta-cyfluthrin Baythroid XL	1.6-2.8 oz	0.013-0.022			7
		beta-cyfluthrin + imidacloprid Leverage 360	3.8-4.1 oz	0.89-0.096			7
		chlorantraniliprole Coragen 1.67 SC	3.5-7.5 fl oz	0.045-0.098			1
		Shenzi 400SC	1.7-3.8 fl oz				
		chlorantraniliprole + lambda-cyhalothrin (R) Voliam Xpress	6-9 oz	0.06-0.09			5
		Cyantraniliprole Exirel 0.83 EC	7-13.5 fl oz	0.45-0.88			1
		lambda-cyhalothrin (R) Karate Z	1.28-1.92 fl oz	0.02-0.03			5
		methomyl (R) Lannate 2.4 LV Lannate 90 SP	1 1/2 pt 1/2 lb	0.45			3
		methoxyfenozide Intrepid 2F	4-16 fl oz	0.06-0.25			1
		zeta-cypermethrin (R) Mustang Maxx 0.8 EC	2.24-4.0 oz	0.014-0.025			1
		zeta-cypermethrin + bifenthrin Hero	4.0-10.3 oz	0.4-0.1			7
POTATOES (white or Irish)	Aphids	beta-cyfluthrin + imidacloprid Leverage 360	2.8 oz	0.066			7
		flonicamid Beleaf 50 SG	2.0-2.8 oz	0.026-0.69			7
		sulfoxaflor Transform WG	0.75-1.0	0.023-0.031			
		thiamethoxam Actara 25 WDG	3 oz	0.047			14
	Colorado potato beetle, potato leafhopper, flea beetles	beta-cyfluthrin Baythroid XL	1.6-2.8 oz	0.013-0.022			0
		beta-cyfluthrin + imidacloprid Leverage 360	2.8 oz	0.066			7

VEGETABLE INSECT CONTROL RECOMMENDATIONS (COMMERCIAL)

Crop	Insect	Insecticide	Formulation Per Acre	Lb ai/Acre	Application	Remarks/Precautions	Days to Harvest
POTATOES (white or Irish) (cont.)	Colorado potato beetle, potato leafhopper, flea beetles (cont.)	carbaryl		0.5-2.0	When insects first appear.	DO NOT apply during bloom. Use higher rates on Colorado potato beetle.	7
		Sevin 4F	1/2-2 qt				
		Sevin XLR Plus	1/2-2 qt				
		esfenvalerate (R)		0.03-0.05			7
		Asana XL 0.66 EC	5.8-9.6 oz				
		lambda-cyhalothrin (R)		0.02-0.03			7
		Karate Z	1.28-1.92 fl oz				
		permethrin (R)				Apply no more than 12 applications. DO NOT graze or feed foliage.	7
		Ambush 25 WP	3.2-12.8 oz	0.05-0.2			
		Ambush 2.0 EC	3.2-12.8 oz	0.05-0.2			
		Pounce 25 WP	6.4-12.8 oz	0.1-0.2			
		spinosad		0.047-0.094			7
		SpinTor 2 SC	3-6 fl oz				
		sulfoxaflor		0.047-0.071			7
		Transform WG	1.5-2.25			For leaf hopper control. Check label for crop rotation restrictions.	
		zeta-cypermethrin (R)		0.02-0.025			1
		Mustang Maxx 0.8 EC	3.2-4.0 oz				
Wireworms		bifenthrin (R)			See label directions.	Check label for crop rotation restrictions.	7
		Capture LFR					
RADISHES	Aphids	phorate (R)			Distribute granules evenly in furrow or band on each side of row at planting or sidedress postemer-gence. Sandy soils use low rate.		90
		Thimet 20G	8.5-17.3 lb				
		beta-cyfluthrin + imidacloprid					
		Leverage 360	2.4-2.8 oz	0.056-0.066			7
	Flea beetles	imidacloprid		0.044			7
		Admire Pro 4.6F	1.3 fl oz				
		malathion		1	When insects appear in damaging numbers.		7
		Malathion 57% EC	1 1/2 pt				
		sulfoxaflor		0.023-0.031		Check label for crop rotation restrictions.	7
		Transform WG	0.75-1.0				
		beta-cyfluthrin		0.013-0.022			0
		Baythroid XL	1.6-2.8 oz				
		beta-cyfluthrin + imidacloprid					
		Leverage 360	2.4-2.8 oz	0.056-0.066			7
		carbaryl		0.5-1.0	When insects appear in damaging numbers.	DO NOT apply during bloom.	3
		Sevin 4F	1/2-1 qt				
		Sevin XLR Plus	1/2-1 qt				
		esfenvalerate (R)		0.03-0.05			7
		Asana XL 0.66 EC	5.8-9.6 oz				
		imidacloprid		0.044			7
		Admire Pro 4.6F	1.3 fl oz				

VEGETABLE INSECT CONTROL RECOMMENDATIONS (COMMERCIAL)

Crop	Insect	Insecticide	Formulation Per Acre	Lb ai/Acre	Application	Remarks/Precautions	Days to Harvest
SOUTHERN PEAS	Bean leaf beetle	beta-cyfluthrin + imidacloprid Leverage 360	2.4-2.8 oz	0.056-0.066			7
		bifenthrin (R) Brigade 2 EC	2.1-6.4 oz	0.033-0.1			3
		carbaryl Sevin 4F Sevin XLR Plus	1/2-1 qt 1/2-1 qt	0.5-1.0	Plant early. Apply treatment to prevent defoliation.	DO NOT apply during bloom. DO NOT use on succulent shelled peas (Group 6B).	3
		lambda-cyhalothrin (R) Karate Z	1.28-1.92 fl oz	0.02-0.03			7
		zeta-cypermethrin (R) Mustang Maxx 0.8 EC	2.24-4.0 oz	0.014-0.025			1
	Corn earworm	beta-cyfluthrin Baythroid XL	1.6-2.1 oz	0.013-0.017			3
		beta-cyfluthrin + imidacloprid Leverage 360	2.4-2.8 oz	0.056-0.066			7
		bifenthrin (R) Brigade 2 EC	2.1-6.4 oz	0.033-0.1			3
		chlorantraniliprole Coragen 1.67 SC Shenzi 400SC	3.5-7.5 fl oz 1.7-3.8 fl oz	0.045-0.098			1
		Cyantraniliprole Exirel 0.83 EC	7-13.5 fl oz	0.45-0.88			1
Cowpea aphid	Cowpea aphid	esfenvalerate (R) Asana XL 0.66 EC	5.8-9.6 oz	0.03-0.05		DO NOT feed vines.	21
		lambda-cyhalothrin (R) Karate Z	1.28-1.92 fl oz	0.02-0.03			7
		methoxyfenozide Intrepid 2F	10-16 oz	0.16-0.25			7
		zeta-cypermethrin (R) Mustang Maxx 0.8 EC	2.24-4.0 oz	0.014-0.025			1
		beta-cyfluthrin + imidacloprid Leverage 360	2.4-2.8 oz	0.056-0.066			7

VEGETABLE INSECT CONTROL RECOMMENDATIONS (COMMERCIAL)

Crop	Insect	Insecticide	Formulation Per Acre	Lb ai/Acre	Application	Remarks/Precautions	Days to Harvest
SOUTHERN PEAS (cont.)	Cowpea aphid (cont.)	malathion Malathion 57% EC 5 lb/gal	1 1/2 pt	1			3
		methomyl (R) Lannate 1.8 LV Lannate SP	1 1/3-3 pt 1/2-1 pt	0.4-0.9		DO NOT apply more than 4.5 lb ai per acre per crop.	1
		zeta-cypermethrin (R) Mustang Maxx 0.8 EC	3.2-4.0 oz	0.02-0.025		Aphid control may be variable.	1
	Cowpea curculio	beta-cyfluthrin Baythroid XL	1.6-2.1 oz	0.013-0.017			3
		beta-cyfluthrin + imidacloprid Leverage 360	2.4-2.8 oz	0.056-0.066			7
		esfenvalerate (R) Asana XL 0.66 EC	5.8-9.6 oz	0.03-0.05			21
		lambda-cyhalothrin (R) Karate Z	1.28-1.92 fl oz	0.02-0.03			7
		zeta-cypermethrin (R) Mustang Maxx 0.8 EC	2.24-4.0 oz	0.014-0.025			1
	European corn borer	beta-cyfluthrin + imidacloprid Leverage 360	2.4-2.8 oz	0.056-0.066			7
		chlorantraniliprole Coragen 1.67 SC Shenzi 400SC	3.5-7.5 fl oz 1.7-3.8 fl oz	0.045-0.098			1
		Cyantraniliprole Exirel 0.83 EC	7-13.5 fl oz	0.45-0.88			1
		methoxyfenozide Intrepid 2F	8-16 oz	0.12-0.25			7
		spinosad SpinTor 2 SC	4-6 fl oz	0.062-0.094			3
SPINACH	Aphids, plant bugs, leafhoppers, leafminers	acetamiprid Assail 70WP	0.8-1.2 oz	0.035-0.054			7
		beta-cyfluthrin + imidacloprid Leverage 360	3 oz	0.07			7
		dimethoate Dimethoate 4 EC	1/2 pt	0.25		Effectiveness of dimethoate is reduced when aphid colonies are established.	14
		dinotefuran Venom 70 SG	1-3 oz foliar 5-6 oz soil	0.044-0.131 0.219-0.263			7 21
		flonicamid Beleaf 50 SG	2.0-2.8 oz	0.062-0.089			0
		imidacloprid Admire Pro 4.6 F	4.4-10.5 fl oz	0.158-0.337		Soil application.	21
		imidacloprid Admire Pro 4.6F	1.3 fl oz	0.047		Imidacloprid should only be used with an aphid scouting program. Apply only when aphid numbers exceed the threshold.	7
		zeta-cypermethrin (R) Mustang Maxx 0.8 EC	3.2-4.0 oz	0.02-0.025			1

VEGETABLE INSECT CONTROL RECOMMENDATIONS (COMMERCIAL)

Crop	Insect	Insecticide	Formulation Per Acre	Lb ai/Acre	Application	Remarks/Precautions	Days to Harvest
SPINACH (cont.)	Caterpillars	<i>Bacillus thuringiensis</i>	Follow label directions.				
		beta-cyfluthrin Baythroid XL	2.4-3.2 oz	0.019-0.025			0
		beta-cyfluthrin + imidacloprid Leverage 360	3 oz	0.07			7
		bifenthrin (R) Brigade 2 EC	2.1-6.4 oz	0.033-0.1		Pyrethroid use against Leps. can result in increases in aphid populations.	40
		chlorantraniliprole Coragen 1.67 SC Shenzi 400SC	3.5-7.5 fl oz 1.7-3.8 fl oz	0.045-0.098			1
		Cyantraniliprole Exirel 0.83 EC	7-13.5 fl oz	0.45-0.88			1
		methomyl (R) Lannate 2.4 LV Lannate SP	1 1/2-3 pt 1/2-1 lb	0.45-0.9	When insects appear or as needed to keep crop free of insects. Not labeled on corn earworm.	DO NOT apply to seedlings less than 3 inches in diameter.	7
		methoxyfenozide Intrepid 2F	4-8 oz	0.06-0.12			1
		permethrin (R) Ambush 2.0 EC Ambush 25 WP Pounce 3.2 EC	6.4-12.8 oz 6.4-12.8 oz 4-8 oz	0.1-0.2			7
		spinosad SpinTor 2 SC	3-6 fl oz	0.047-0.094			1
		thiodicarb (R) Larvin 3.2	24-30 oz	0.6-0.75			14
		zeta-cypermethrin (R) Mustang Maxx 0.8 EC	3.2-4.0 oz	0.02-0.025			1
SWEET CORN	Corn Earworm	beta-cyfluthrin Baythroid XL	1.6-2.8 oz	0.013-0.022			0
		bifenthrin (R) Brigade 2 EC	2.1-6.4 fl oz	0.033-0.1	When larvae are found.		1
		carbaryl Sevin 4F Sevin XLR	1-2 qt 1-2 qt	1.0-2.0		DO NOT apply during bloom.	2
		chlorantraniliprole Coragen 1.67 SC Vantacor Shenzi 400SC	3.5-7.5 fl oz 1.2-2.5 fl oz 1.7-3.8 fl oz	0.045-0.098		Cheaper option than Coragen in Sweet Corn	1
		esfenvalerate (R) Asana XL 0.66 EC	5.8-9.6 oz	0.03-0.05			1
		lambda-cyhalothrin (R) Karate Z	1.28-1.92 fl oz	0.02-0.03			1

VEGETABLE INSECT CONTROL RECOMMENDATIONS (COMMERCIAL)

Crop	Insect	Insecticide	Formulation Per Acre	Lb ai/Acre	Application	Remarks/Precautions	Days to Harvest
SWEET CORN (cont.)	Corn Earworm (cont.)	methomyl (R) Lannate 2.4 LV Lannate SP	1-1 1/2 pt 1/3-1/2 lb	0.3-0.45			0 (ears) 3 (forage)
		methoxyfenozide + spinetoram Intrepid Edge		8-12 oz		Follow label directions.	21 (dry fodder) 3 (forage) 3 (ears)
		permethrin (R) Ambush 2.0 EC Ambush 25 WP		6.4-12.8 oz 6.4-12.8 oz	0.1-0.2		1
		spinosad SpinTor 2 SC		3-6 fl oz	0.047-0.094		1
		thiodicarb (R) Larvin 3.2		20-30 oz	0.5-0.75		0
		zeta-cypermethrin (R) Mustang Maxx 0.8 EC		2.24-4.0 oz	0.014-0.025		3
Cutworms, rootworms, seed corn maggot, white grubs, wireworms	beta-cyfluthrin Baythroid XL		0.8-1.6 oz	0.007-0.013		Cutworms only.	0
	bifenthrin (R) Capture LFR		0.39-0.44 oz/ 1,000 ft row	0.08-0.1	Apply in 5- to 7-inch band or in furrow ahead of press wheel. Compatible with liquid fertilizers.		3
	tefluthrin (R) Force 3G		4-5 oz/ 1,000 ft row	0.075-0.165	Banded, T-banded or in furrow.		
	zeta-cypermethrin + bifenthrin Hero		2.6-6.1 oz	0.025-0.05			3
European corn borer, southwestern corn borer	carbaryl Sevin 4F Sevin XLR Plus		1 1/2-2 qt 1 1/2-2 qt	1.5-2.0	When 50% of plants show rows of tiny pinholes in leaves (feeding signs).	Use high volume sprays directed into whorls. DO NOT apply during bloom.	0
	chlorantraniliprole Coragen 1.67 SC Shenzi 400SC		3.5-7.5 fl oz 1.7-3.8 fl oz	0.045-0.098			1
	esfenvalerate (R) Asana XL 0.66 EC		5.8-9.6 oz	0.03-0.05			1
	methomyl (R) Lannate 2.4 LV Lannate SP		1 1/2 pt 1/2 lb	0.45		Lannate not labeled on southwestern corn borer.	0 (ears) 3 (forage)
	methoxyfenozide Intrepid 2F		4-16 oz	0.06-0.25			21 (dry fodder) 3 (forage) 3 (ears)
	methoxyfenozide + spinetoram Intrepid Edge		4-12 oz				21 (dry fodder) 3 (forage) 3 (ears)

VEGETABLE INSECT CONTROL RECOMMENDATIONS (COMMERCIAL)

Crop	Insect	Insecticide	Formulation Per Acre	Lb ai/Acre	Application	Remarks/Precautions	Days to Harvest
SWEET CORN (cont.)	European corn borer, southwestern corn borer (cont.)	spinosad SpinTor 2 SC	1.5-6.0 fl oz	0.023-0.094			1
		thiodicarb (R) Larvin 3.2	20-30 oz	0.05-0.75			0
	Flea beetles	acetamiprid + bifenthrin Savoy	9.6-12.9 oz			Maximum of 25.8 oz a season.	7
		beta-cyfluthrin Baythroid XL	0.8-1.6 oz	0.007-0.013			0
		carbaryl Sevin 4F Sevin XLR Plus	1-2 qt 1-2 qt	1.0-2.0	When damage appears.	DO NOT apply during bloom. Hand harvesting is prohibited.	2
		esfenvalerate (R) Asana XL 0.66 EC	5.8-9.6 oz	0.03-0.05			1
		zeta-cypermethrin (R) Mustang Maxx 0.8 EC	2.8-4.0 oz	0.017-0.025			3
	Mites	abamectin Agri-Mek SC	1.75-3.5 fl oz	0.01-0.05	For control of spider mites.	Must be mixed with an adjuvant (NIS surfactant).	7
	Stink Bugs	cetamiprid + bifenthrin Savoy	12.9 oz			Maximum of 25.8 oz a season.	7
		bifenthrin (R) Brigade 2 EC	2.1-6.4 oz	0.033-0.1		Follow label directions.	1
		methomyl (R) Lannate 2.4 LV Lannate SP	1/2 lb 1/2 lb	0.45			0 (ears) 3 (forage)
Whorl feeders, fall armyworm, corn earworm	beta-cyfluthrin Baythroid XL	1.6-2.8 oz	0.013-0.022				0
	bifenthrin (R) Brigade 2 EC	2.1-6.4 oz	0.033-0.1			Follow label directions.	1
	carbaryl Sevin 4F Sevin XLR Plus	1-2 qt 1-2 qt	1.0-2.0	Apply when needed. Hand harvesting is prohibited.		Plants will tolerate a great deal of feeding with no yield loss. DO NOT apply during bloom.	2
	chlorantraniliprole Coragen 1.67 SC Shenzi 400SC	3.5-7.5 fl oz 1.7-3.8 fl oz	0.045-0.098				1
	esfenvalerate (R) Asana XL 0.66 EC	5.8-9.6 oz	0.03-0.05				1
	lambda-cyhalothrin (R) Karate Z	1.28-1.92 fl oz	0.02-0.03				1
	methomyl (R) Lannate 2.4 LV Lannate SP	1-1 1/2 pt 1/3-1/2 pt	0.03-0.45				3

VEGETABLE INSECT CONTROL RECOMMENDATIONS (COMMERCIAL)

Crop	Insect	Insecticide	Formulation Per Acre	Lb ai/Acre	Application	Remarks/Precautions	Days to Harvest
SWEET CORN (cont.)	Whorl feeders, fall armyworm, corn earworm (cont.)	spinosad SpinTor 2 SC	3-6 oz	0.042-0.094		Follow label directions.	1
		zeta-cypermethrin (R) Mustang Maxx 0.8 EC	3.2-4.0 oz	0.02-0.025			3
		zeta-cypermethrin + bifenthrin Hero	4.0-10.3 oz	0.04-0.1			3
SWEET POTATO	Flea beetles, tortoise beetles	beta-cyfluthrin Baythroid XL	1.6-2.8 oz	0.013-0.022			0
		beta-cyfluthrin + imidacloprid Leverage 360	2.4-2.8 oz	0.056-0.066			7
		bifenthrin (R) Brigade 2 EC	2.1-6.4 fl oz	0.033-0.1		Follow label directions.	21
		carbaryl Sevin 4F Sevin XLR Plus	1-2 qt 1-2 qt	1.0-2.0	When damage occurs on leaves as needed.	DO NOT apply during bloom.	7
		lambda-cyhalothrin (R) Karate Z	1.28-1.92 fl oz	0.02-0.03			7
		spinosad SpinTor 2 SC	3-6 fl oz	0.047-0.094		Not labeled for tortoise beetle.	7
Sweet potato weevil		beta-cyfluthrin Baythroid XL	1.6-2.8 oz	0.013-0.022			0
		beta-cyfluthrin + imidacloprid Leverage 360	2.4-2.8 oz	0.056-0.066			7
		carbaryl Sevin 4F Sevin XLR Plus	2.6 oz/gal 2.6 oz/gal		Use as a preplant dip for sweet potato weevil. Just prior to plant- ing, dip sweet potato cuttings in the suspension.	DO NOT apply during bloom.	7
		lambda-cyhalothrin (R) Karate Z	1.28-1.92 fl oz	0.02-0.03			7
		phosmet Imidan 5% dust	2-4 oz/50 lb bu stored		One application.		
Wireworm, other soil insects	bifenthrin (R) Capture LFR				See label directions.		

VEGETABLE INSECT CONTROL RECOMMENDATIONS (COMMERCIAL)

Crop	Insect	Insecticide	Formulation Per Acre	Lb ai/Acre	Application	Remarks/Precautions	Days to Harvest
TOMATO	Cabbage loopers	<i>Bacillus thuringiensis</i>			Check label.		
		Dipel					
		Javelin					
		Biobit					
		Agree					
		chlorantraniliprole			0.045-0.098		1
		Coragen 1.67 SC	3.5-7.5 fl oz				
		Shenzi 400SC	1.7-3.8 fl oz				
		Cyantraniliprole			0.45-0.88		1
		Exirel 0.83 EC	7-13.5 fl oz				
Colorado potato beetles, flea beetles, stink bugs	Colorado potato beetles, flea beetles, stink bugs	chlorantraniliprole + lambda-cyhalothrin (R)			0.05-0.08		5
		Voliam Xpress	5-8 oz				
		esfenvalerate (R)			0.03-0.05		1
		Asana XL 0.66 EC	5.8-9.6 oz				
		methomyl (R)			0.45-0.9	Apply on foliage as needed when loopers are small.	1
		Lannate 2.4 LV	1 1/2-3 pt				
		Lannate SP	1/2-1 lb				
		methoxyfenozide			0.06-0.12		1
		Intrepid 2F	4-8 oz				
		methoxyfenozide + spinetoram				Follow label directions.	1
Cutworms	Cutworms	Intrepid Edge	10-12 oz				
		beta-cyfluthrin			2.8 oz for flea beetles.		0
		Baythroid XL	1.6-2.8 oz				
		beta-cyfluthrin + imidacloprid					
		Leverage 360	3.8-4.1 oz		0.089-0.096		0
		bifenthrin (R)			0.033-0.08		1
		Brigade 2 EC	2.1-5.2 oz				
		carbaryl			1.0-2.0	Apply on foliage as needed.	3
		Sevin 4F	1-2 qt				
		Sevin XLR Plus	1-2 qt				
Vegetables (Commercial)	Vegetables (Commercial)	imidacloprid			0.047		0
		Admire Pro 4.6F	1.3 fl oz				
		spinosad			0.047-0.094	Follow label directions.	1
		SpinTor 2 SC	3-6 oz				
		zeta-cypermethrin (R)			0.02-0.025		1
		Mustang Maxx 0.8 EC	3.2-4.0 oz				
		zeta-cypermethrin + bifenthrin					
		Hero	4.0-10.3 oz		0.04-0.1		1
		beta-cyfluthrin + imidacloprid					0
		Leverage 360	3.8-4.1 oz		0.089-0.096		

VEGETABLE INSECT CONTROL RECOMMENDATIONS (COMMERCIAL)

Crop	Insect	Insecticide	Formulation Per Acre	Lb ai/Acre	Application	Remarks/Precautions	Days to Harvest
TOMATO (cont.)	Cutworms (cont.)	bifenthrin (R) Capture LFR	0.29-0.39 oz/ 1,000 ft row OR 3.4-6.8 oz/acre	0.04-0.08	Apply in 5- to 7-inch band or in furrow ahead of press wheel. Compatible with liquid fertilizers.		30
		carbaryl Sevin 4F Sevin XLR Plus	2 qt 2 qt	2.0	Apply to soil surface 2-3 days prior to transplanting or when damage occurs.	DO NOT apply during bloom.	3
		diazinon Diazinon AG500 Diazinon 50 W Diazinon AG600C	2-4 qt 4-8 lb 51-102 oz	2.0-4.0	Broadcast to soil surface just prior to planting. Incorporate into soil 3-6 inches immediately. Mix in a minimum of 5 gallons of water/acre.		
		esfenvalerate (R) Asana XL 0.66 EC	5.8-9.6 oz	0.03-0.05	Broadcast prior to planting and work into soil 4-8 inches.		1
		zeta-cypermethrin (R) Mustang Maxx 0.8 EC	1.28-4.0 oz	0.008-0.025			1
		zeta-cypermethrin + bifenthrin Hero	10.3 oz	0.1			1
Leaf miners, aphids	acetamiprid + bifenthrin Savoy		4.9-9.6 oz			Aphids only. Follow label directions.	7
	abamectin (R) Agri-Mek 0.7 SC		1.75-3.5 fl oz	0.009-0.019	20 gallons of water/acre required.	Mites, leafminers, pinworms, potato beetle. Must be mixed with an adjuvant (NIS surfactant).	7
	acetamiprid Assail 30 SG		2-4 oz	0.038-0.075		Follow label directions.	7
	beta-cyfluthrin + imidacloprid Leverage 360		3.8-4.1 oz	0.089-0.096			0
	bifenthrin (R) Brigade 2 EC		2.1-5.2 oz	0.033-0.08			1
	dimethoate Dimethoate 400 Dimethoate 4EC Dimethoate 5 lb			0.25-0.5			7
	esfenvalerate (R) Asana XL 0.66 EC		9.6 oz	0.05		Use 5.8-9.6 oz of Asana per acre for aphids.	1
	imidacloprid Admire Pro 4.6F		1.3 fl oz	0.047			0
	methomyl (R) Lannate 2.4 LV Lannate SP		1 1/2-3 pt 1/2-1 lb	0.45-0.9		Aphids only.	1
	sulfoxaflor Transform WG		0.75-1.0	0.023-0.031		Not for leaf miners. Check label for crop rotation restrictions.	1
	zeta-cypermethrin (R) Mustang Maxx 0.8 EC		3.2-4.0 oz	0.02-0.025		Aphid control may be variable.	1
	zeta-cypermethrin + bifenthrin Hero		4.0-10.3 oz	0.04-0.1			1

VEGETABLE INSECT CONTROL RECOMMENDATIONS (COMMERCIAL)

Crop	Insect	Insecticide	Formulation Per Acre	Lb ai/Acre	Application	Remarks/Precautions	Days to Harvest
TOMATO (cont.)	Pinworms	abamectin (R) Agri-Mek 0.7 SC	3.5 oz	0.019		Must be mixed with an adjuvant (NIS surfactant).	7
		beta-cyfluthrin Baythroid XL	2.1-2.8 oz	0.017-0.022			0
		beta-cyfluthrin + imidacloprid Leverage 360	3.8-4.1 oz	0.089-0.096			0
		chlorantraniliprole Coragen	3.5-7.5 fl oz	0.045-0.098			1
		esfenvalerate (R) Asana XL 0.66 EC	5.8-9.6 oz	0.03-0.05		DO NOT exceed 0.5 lb ai/acre per season.	1
		gamma-cyhalothrin Proaxis 0.5 EC	2.56-3.84 oz	0.01-0.015		Follow label directions.	5
		indoxacarb Avaunt 30 DG	3.5 oz	0.065		Follow label directions.	3
		methomyl (R) Lannate LV Lannate SP	1 1/2-3 pt 1/2-1 lb	0.45-0.9		All are alternatives. Check dosage on label.	1
		methoxyfenozide Intrepid 2F	10-16 oz	0.16-0.25			1
		No-Mate Spiral Bioinsecticide			Read product label.	Comes in a plastic coil that is wrapped around tomato vines or support stakes.	
Spider mites		abamectin (R) Agri-Mek 0.7 SC	1.75-3.5 oz	0.0094-0.019	DO NOT apply Agri-Mek in less than 20 gallons of water per acre.	Must be mixed with an adjuvant (NIS surfactant).	7
		bifenthrin (R) Brigade 2 EC	5.2-6.4 oz	0.08-0.1		Follow label directions.	1
		fenpyroximate Portal	2 pt		Apply in a minimum of 20 gallons of water per acre		1
		spiromesifen Oberon 2 SC	7-8.5 oz	0.11-0.13			1
Thrips		acetamiprid + bifenthrin Savoy	9.6 oz			Follow label directions.	7
		beta-cyfluthrin + imidacloprid Leverage 360	3.8-4.1 oz	0.089-0.096			0
		bifenthrin (R) Brigade 2 EC	2.1-5.2 oz	0.033-0.08			1
		flonicamid Beleaf	2.4-4.8 oz	0.089-0.133			0
		spinetoram Radiant 1 SC	6-10 oz	0.047-0.078		Will control thrips on foliage, but not in flowers.	1

VEGETABLE INSECT CONTROL RECOMMENDATIONS (COMMERCIAL)

Crop	Insect	Insecticide	Formulation Per Acre	Lb ai/Acre	Application	Remarks/Precautions	Days to Harvest
TOMATO (cont.)	Tomato fruitworm, Hornworms	carbaryl Sevin 4F Sevin XLR Plus	1-2 qt 1-2 qt	1.0-2.0		DO NOT apply during bloom.	3
		chlorantraniliprole Coragen 1.67 SC Shenzi 400SC	3.5-7.5 fl oz 1.0-3.8 fl oz	0.045-0.098		2.0-5.0 fl oz for hornworms	1
		Cyantraniliprole Exirel 0.83 EC	7-13.5 fl oz	0.45-0.88			1
		chlorantraniliprole + lambda-cyhalothrin (R) Voliam Xpress	6-9 oz	0.06-0.09			5
		esfenvalerate (R) Asana XL 0.66 EC	5.8-9.6 oz	0.03-0.05			1
		lambda-cyhalothrin (R) Karate Z	1.28-1.92 fl oz	0.02-0.03	Five-day intervals.		5
		methomyl (R) Lannate 2.4 LV Lannate SP	1 1/2-3 pt 1/2-1 lb	0.45-0.9		<i>Bacillus thuringiensis</i> 1/2 lb/acre may be used in combination with methomyl. Other areas have had good results.	
		methoxyfenozide Intrepid 2F	4-16 oz	0.06-0.25			1
		methoxyfenozide + spinetoram Intrepid Edge	10-12 oz	0.06-0.25		Follow label directions.	1
		rynavypyr Coragen	3.5-5 fl oz	0.045-0.065			1
		spinosad SpinTor 2 SC	3-8 fl oz	0.047-0.125			1
		zeta-cypermethrin (R) Mustang Maxx 0.8 EC	2.24-4.0 oz	0.014-0.025			1
		zeta-cypermethrin + bifenthrin Hero	4.0-10.3 oz	0.04-0.1			1
	Wireworms	bifenthrin (R) Capture LFR	0.29-0.39 oz/ 1,000 ft row or 3.4-6.8 oz/A	0.04-0.08	Apply in 5- to 7-inch band or in furrow ahead of press wheel. Compatible with liquid fertilizers.		30
		diazinon (R) Diazinon 50W Diazinon AG600 Diazinon AG500	6-8 lb 76 1/2-102 oz 3-4 qt	3.0-4.0	Broadcast prior to planting and immediately work into soil 4-8 inches.		
VEGETABLES (See labels for specific crops)	Red Imported Fire Ants	ALL FIRE ANT BAITS – Apply when ants are active and soil temperature is above 60 degrees F. DO NOT treat if rain is anticipated within 6 hours.					
		S-Methoprene Extinguish	3-5 Tbsp/mound 1.0-1.5 lb/acre		Mound-to-mound treatment rate. Broadcast rate.	This product is an insect growth regulator (IGR). Apply 3 to 4 weeks prior to harvest date.	
		pyriproxyfen Esteem	2-4 Tbsp/mound 1.5-2.0 lb/acre		Mound-to-mound treatment rate. Broadcast rate.	This product is an insect growth regulator (IGR).	
		spinosad Eliminator Fire Ant Bait Green Light Fire Ant Control Ortho Fire Ant Bait Safer Fire Ant Bait	4-6 Tbsp/mound 2.5-5.0 lb/acre		Mound-to-mound treatment rate. Broadcast rate.	See label for specific rate and application instructions. Acceptable for organic production.	