

2020 Facts:

- 12,000 acres planted (estimated)
- 77 bu/acre average yield (estimated)
- Key dates, 2007-2018 Verification Program
 Click here for more information about verification

Planting Date: April 22
 Emergence: April 29
 Harvest: August 30

• 56 lbs = 1 bu

14% moisture is dry

Growth and Development:

	Average from GSRVP		
	2007-2018		
	Days	Inches	Applications
Planting	0	0	~ April 22
VE	7	0	
V1	10	1	
V2	13	2	
V3	17	3	
V4	22	4	Cidaduasa
V5	26	5	Sidedress Fertilizer
V6	30	7	rerunzer
V7	35	9	Atrazine Cutoff (12 in)
V8	39	12	
Boot	66	36	Scout for Foliar
Heading	73	49	Disease
Flowering	76	53	Scout for Midge
Soft Dough	86	53	Coout for
Hard	0.7	F2	Scout for Headworms
Dough	97	53	пеаимогті
Maturity	112	53	
Harvest	131	53	~ August 30

Seeding:

- Seed should be Concep treated to allow the use of metolachlor (Dual) herbicide
- Place seed between 0.75 to 1.5 inches deep

- Plant when ground temp is 65° @ 2 inches deep by 9:00 a.m. for three days
- Irrigated seeding rate should be 80,000 -100,000 seeds per acre with a final plant population of 75,000 plants per acre assuming 80% germination
- Dryland seeding rate should be 75,000 seeds per acre with a final plant population of 50,000 - 60,000 plants per acre assuming 80% germination
- Consider an insecticide seed treatment such as Cruiser, Poncho, etc. for improving stands

Grain Sorghum Seeding Rates

_		_		
Seeding	Row Spacing (inches)			
Rate	7"	30"	38"	Final Stand
(seeds				at 80%
per	Seeds	s per 10 f	t row	Germination
acre)				
55,000	7.4	31.6	40.0	44,000
60,000	8.0	34.4	43.6	48,000
65,000	8.7	37.3	47.3	52,000
70,000	9.4	40.2	50.9	56,000
75,000	10.0	43.0	54.5	60,000
80,000	10.7	45.9	58.2	64,000
85,000	11.4	48.8	61.8	68,000
90,000	12.1	51.7	65.4	72,000
95,000	12.7	54.5	69.1	76,000
100,000	13.4	57.4	72.7	80,000

Determining Final Plant Populations:

- 30" rows measure 17 ft 5 in
- 38" rows measure 13 ft 9 in

Count plants in that distance and multiply by 1,000. This will equal plants per acre. Do this in at least ten stops in the field to get an accurate count. Example: 30" row, count 80 plants in 17 ft 5 in

80 X 1000 = 80,000 plants per acre

Seeding Rate in Pounds per acre

Seeds per	Seeds per Pound				
Acre	11,000	13,000	15,000	17,000	
7.0.0	Pounds of Seed per Acre				
60,000	5.5	4.6	4.0	3.5	
70,000	6.4	5.4	4.7	4.1	
80,000	7.3	6.2	5.3	4.7	
90,000	8.2	6.9	6.0	5.3	
100,000	9.1	7.7	6.7	5.9	
110,000	10.0	8.5	7.3	6.3	

Hybrid Selection:

2020 Yields - Arkansas Grain Sorghum Hybrid Testing Program

	Irrigated Yield	Non Irr. Yield
Hybrid	(bu/ac)	(bu/ac)
Dekalb DKS 53-53	153	135
Dyna-Gro 69BG38	153	123
Dyna-Gro M71GR91*	152	131
Dekalb DKS 51-01	150	121
Dyna-Gro M72GB71*	147	120
Dekalb DKS 45-23	147	124
Sorghum P. 7715*	147	111

^{*}Sugarcane Aphid Tolerant

More information on Arkansas Grain Sorghum Hybrid Trials can be found at:

https://aaes.uark.edu/variety-testing/

Fertilization:

Nitrogen (N):

- Apply ¼ to ⅓ of N immediately before or after planting
- Apply sidedress N between the V4 and V6 stage

N Recommendation

Yield Goal	Units of N to apply per acre		
(bu/ac)	Sandy, Silt Loams	Clay	
Dryland	110	150	
90	110	150	
110	135	175	
<u>≥</u> 130	160	200	

Nitrogen sources:

- 32% UAN (1 gal = 3.5 lbs of N)
- Urea (46-0-0)
- DAP (18-46-0)
- Ammonium Sulfate (21-0-0-24)

Phosphorus (P) and Potassium (K):

P₂O₅ Recommendation

Yield	Soil Test P (ppm)			
Goal	<16	16-25	26-35	>36
(bu/ac)	lbs of P ₂ O ₅ per acre			
Dryland	90	70	50	0
90	90	70	50	0
110	100	80	60	0
<u>></u> 130	110	90	70	0

K₂O Recommendation

Yield		Soi	l Test K (ppm)	
Goal	<61	61-	91-	131-	>175
	<01	90	130	175	21/3
(bu/ac)		lbs c	of K₂O pe	er acre	
Dryland	140	90	60	0	0
90	140	90	60	0	0
110	140	100	70	0	0
<u>></u> 130	150	100	70	40	0

Diseases and Fungicide Timing:

- Fungicides should only be applied when disease is present
- Plant resistant hybrids to help with anthracnose and target leaf spot

 Check the MP 154 for the latest fungicide recommendations (click for electronic copy)

Irrigation:

Potential Yield Reduction from Moisture Stress		
Growth Stage % Yield Reduction		
Emergence to V8	10 – 15	
Boot to Flowering	30 – 50	
Soft dough to maturity	10 – 20	

Estimated Grain Sorghum Water Use		
Days after planting	Inches/day	
0-30 (early plant growth)	0.05 - 0.10	
30-60 (rapid plant growth)	0.10 - 0.20	
60-80 (boot and flowering)	0.25 - 0.30	
80-120 (grain fill to maturity)	0.25 - 0.10	

Irrigation Termination

- Furrow Irrigation –when >50% of the heads are at hard dough and there is adequate moisture
- Pivot Irrigation –when >75% of the heads are at hard dough and there is adequate moisture

Herbicides:

- Make sure seed is treated with Concep prior to planting
- Apply metolachlor (Dual Magnum) PRE followed by atrazine plus Dual Magnum POST
- 1 qt of atrazine 4L = 1 lb of atrazine
- Do not apply more than 2.5 lb of atrazine in a season
- Do not apply atrazine after 12 inches
- Apply atrazine to healthy and growing grain sorghum to avoid injury

- There is a 10 month plant back on grain sorghum following fomesafen (Prefix, Flexstar)
- There is a 18 month plantback following Newpath
- Check the MP 44 for the latest herbicide recommendations (click for electronic copy)

Insects:

 Check the MP 144 for the latest insecticide recommendations (click for electronic copy)

Insect	Threshold
Whorl Feeders	3-6 larvae per whorl
Chinch Bug	When stand is threatened
Greenbug	Plants up to 6 inches – visible damage 6 inches to preboot – treat before entire leaves die Preboot to hard dough – treat when 2 or more entire leaves die
Sorghum Midge	1 adult midge per head when 25% - 30% of plants are flowering
Sugarcane Aphid	25% of plants infested with a colony of at least 50 aphids
Headworm Complex (Corn Earworm + Fall Armyworm)	1 larvae ½ inch long or longer per head
Sorghum Webworm	5 to 6 larvae per head
Stink Bug	5 per head from flowering to soft dough 16 per head during hard dough

More information and additional copies of this fact sheet are available at:

www.uaex.uada.edu/grain-sorghum www.uaex.uada.edu/verification