Dr. Jason Kelley – Extension Agronomist – Wheat and Feed Grains Chuck Capps – Program Associate – Corn and Grain Sorghum Verification



2023 Facts:

- 850,000 acres planted
- 183 bushel/acre state average yield
- Average dates in 2023 CRVP
 <u>Click here for more information about verification</u>
 - Planting: April 10
 - o Emergence: April 19
 - Harvest: September 5
- 56 lbs = 1 bu
- 15.5% moisture is dry

Corn Growth and Development:

	Average from Verification Program			
	Heat			
	Days	Inch	Unit	Applications
Plant	0	0		April 10
VE	9	0	147	
V2	18	3	279	
V4	27	6	442	
V6	36	12	612	Atrazine & Permit Cutoff (12 in)
V8	44	21	794	Callisto Glyphosate Cutoff (30 in)
V10	50	35	947	
V12	55	51	1083	Pre-Tassel N
V14	60	65	1194	Pre-Tasser N
V16	64	80	1306	
R1	71	116	1497	
R2	77	116	1665	Fungicide
R3	83	116	1847	
R4	90	116	2036	
R5	97	116	2230	Irrigation
R6	121	116	2870	Termination
Harvest	148	116		Sept 5

Seeding:

• Plant when soil temperature is 55° @ 2 inches deep by 9:00 a.m. for three days

- Plant seed 2 inches deep in most instances
- Seeding rates for irrigated fields, 32,000 to 34,000 seeds/acre, 25,000 for non-irrigated fields.
- Avoid planting 48 hours or less ahead of cold rains if possible to ensure maximum emergence
- Greatest yields are generally obtained from corn planted before April 25th in South Arkansas and May 1 in North Arkansas

Corn Seeding Rates

Row Spacing (inches)			
30″	38"	30″	38"
See	eds	Inc	hes
per 10 Feet of Row		Between Seed	
14.9	18.9	8.0	6.3
16.1	20.4	7.5	5.8
17.2	21.8	7.0	5.5
18.4	23.3	6.5	5.2
19.5	24.7	6.1	4.9
20.7	26.2	5.8	4.6
21.8	27.6	5.5	4.3
23.0	29.1	5.2	4.1
	30" See per 10 Fe 14.9 16.1 17.2 18.4 19.5 20.7 21.8	30" 38" Seeds per 10 Feet of Row 14.9 18.9 16.1 20.4 17.2 21.8 18.4 23.3 19.5 24.7 20.7 26.2 21.8 27.6	$\begin{array}{c c c c c c c c c c c c c c c c c c c $

Determining growth stage:

- Corn growth stages are designated V for vegetative and R for reproductive
- Each V number represents the upper most leaf with a visible collar (Ex:V2 = 2 leaf)
- A corn plant typically has 19 21 leaves

Vegetative Stages		Reproductive Stages		
VE	emergence	R1	silking	
V1	1 leaf	R2	blister	
V2	2 leaf	R3	milk	
V3	3 leaf	R4	dough	
V(n)	n th leaf	R5	dent	
VT	tasseling	R6	physiological maturity	

Determining Final Plant Populations:

- 30" rows measure 17 ft 5 in
- 38" rows measure 13 ft 9 in

Count plants in that distance and multiply by 1,000. This will equal plants per acre. Do this in at least ten stops in the field to get an accurate count. Example: 30" row, count 34 plants in 17 ft 5 in 34 X 1000 = 34,000 plants/acre

Fertilization:

Nitrogen (N):

- Apply approximately ¼ to ¼ of N immediately before or immediately after planting
- Apply sidedress N between the V4 and V6 growth stages
- If applying pretassel N, apply 100 lbs/acre of Urea (46 units) one to two weeks prior to tassel (approximately between V12 and V14 stage)

Yield				
Goal	Units of N to apply/acre			
(bu/ac)	Sandy, Silt Loams	Clay		
125	160	230		
150	160	230		
175	220	290		
<u>></u> 200	220	290		

Nitrogen sources:

- 32% UAN (1 gal = 3.5 units of N)
- Urea (46-0-0)
- DAP (18-46-0)
- Ammonium Sulfate (21-0-0-24)
- Ammonium Polyphosphate (10-34-0)

Phosphorus (P) and Potassium (K):

P_2O_5 Recommendation

Yield	Soil Test P (ppm)				
Goal	<u><</u> 8	9-16	17-35	36-50	<u>></u> 51
(bu/ac)	lbs of P ₂ O ₅ per acre				
125	80	60	40	0	0
150	100	70	50	0	0
175	120	80	60	0	0
<u>></u> 200	120	90	70	0	0

K₂O Recommendation

Vi a lal	Soil Test K (ppm)				
Yield	-61	61.00	91-	131-	► 17 Γ
Goal	<61 61-90	61-90	130	175	>175
(bu/ac)	lbs of K ₂ O per acre				-
125	100	80	60	0	0
150	120	100	60	0	0
175	150	110	60	0	0
> 200	160	120	70	50	0

Zinc (Zn):

- Apply 10 lbs/ac of Zn as a granular when Zn levels are <4 ppm and pH is >6.0
- 33 lbs/ac of Zinc Sulfate applied preplant equals approximately 10 lbs/ac of actual Zn

Sulfur (S):

- Apply 20 lbs/ac of S when the SO₄-S soil test level is <10 ppm or a deficiency has occurred in the past
- 100 lbs/ac of Ammonium Sulfate equals 24 lbs of actual S

Diseases and Fungicide Timing:

- Fungicides should only be applied when disease is present
- Silk to brown silk and later is when we typically see foliar disease develop
- Common rust has a brick-red color with circular to elongated pustules, it comes in

early and usually does not require treatment

- Southern rust has an orange pustule and comes in later in the year, the pustules are usually on the upper leaf surface and can require a fungicide
- <u>Check the MP 154 for the latest fungicide</u> recommendations (click for electronic copy)

Irrigation:

Potential Yield Reduction from Moisture Stress		
Growth Stage	% Yield Reduction	
Prior to tasseling	10 - 20	
Tasseling to soft dough	20 – 60	
Soft dough to maturity	10 - 35	

Estimated Corn Water Use*			
Days after planting	Inches/day		
0-30 (early plant growth)	0.05 - 0.10		
30-60 (rapid plant growth)	0.10 - 0.20		
60-100 (reproductive stage)	0.20 - 0.30		
100-120 (grain fill to maturity) 0.25 – 0.10			
* Based on planting date of April 1			

Irrigation Termination

- Furrow Irrigation when the starch line on representative kernels has moved >50%, and there is adequate moisture
- Pivot Irrigation when the starch line on representative kernels has moved >75%, and there is adequate moisture

Herbicides:

- 1 qt of atrazine 4L = 1 lb of atrazine
- Do not apply >2.5 lb/ac of atrazine in a season
- For best weed control, apply metolachlor (Dual II Magnum) or other residual PRE

followed by POST herbicide combination including atrazine by V5

- Do not apply atrazine after corn more than 12 inches tall
- Halex GT, Callisto, and glyphosate can be applied up to 30-inch corn or V8 growth stage, whichever is more restrictive
- <u>Check the MP 44 for the latest herbicide</u> recommendations (click for electronic copy)

Insects Traits:

Bt traits commonly utilized in Arkansas Corn

		Insects	
Trait	Symbol	Managed	Refuge
Genuity VT Double Pro	DPRO	Corn Borers Fall Armyworm	20%
Genuity VT Triple Pro	PRO	Corn Borers Fall Armyworm Corn Rootworm	20%
Genuity SmartStax	SS	Corn Borers Fall Armyworm Corn Rootworm	20%
Genuity Tricepta	TRE	Corn Borers, Fall Armyworm, Corn Earworm	20%
Agrisure Viptera	VIP	Corn Borers Fall Armyworm Corn Earworm	20%
Optimum AcreMax Leptra	VYHR	Corn Borers Fall Armyworm Corn Earworm	20%
Optimum AcreMax	YHR	Corn Borers Fall Armyworm	20%

For more information:

https://www.uaex.uada.edu/farm-ranch/cropscommercial-horticulture/corn/

The University of Arkansas System Division of Agriculture offers all its Extension and Research programs and services without regard to race, color, sex, gender identity, sexual orientation, national origin, religion, age, disability, marital or veteran status, genetic information, or any other legally protected status, and is an Affirmative Action/Equal Opportunity Employer.