

## Livestock Health Series

# Internal Parasites in Beef and Dairy Cattle

Shane Gadberry  
Associate Professor -  
Animal Science

Jeremy Powell  
Associate Professor -  
Veterinarian

The need to control internal parasites will exist as long as cattle are grazing pastures. However, parasite levels are not the same on all pastures or in all cattle. Pastures that are heavily stocked generally have a higher parasite burden than lightly stocked ones. Cattle in a drylot are less likely to have heavy worm infections than those on pastures. Young cattle will typically have more internal parasites than older cattle. Therefore, the methods of controlling internal parasites should be developed to fit individual production situations. Strategic deworming starts with understanding the life cycle of problem parasites, identifying seasonal changes in parasite burdens and implementing cost-effective control. A successful deworming program, along with good overall herd management, will increase milk production in cows and thereby increase weaning weights of calves.

### Effects of Internal Parasites

The effects of internal parasites on cattle will vary with the severity of infection as well as age and stress level of the animal. In general, younger animals and animals under stress are most likely to show signs of parasitism. Mature cows acquire a degree of immunity to parasites that reside in the lower gastrointestinal tract. However, the brown stomach worm (*Ostertagia ostertagi*) has evolved to evade the animal's immune system. In addition, parasite burdens are most detrimental in mature cows

near parturition because immunity is suppressed. Cows, especially dairy, in early lactation are often in a negative energy balance due to the stress of lactation. These cattle are affected more than cows in later lactation, when smaller levels of milk are being produced. Bulls are typically more susceptible to internal parasites than cows.

The effects of parasitism can be separated into two types – **subclinical** and **clinical**. Losses in animal productivity (milk production, weight gain, altered carcass composition, conception rate, etc.) are all subclinical effects; whereas, visible, disease-like symptoms (roughness of coat, anemia, edema, diarrhea) are clinical effects. The subclinical effects are of major economic importance to the producer.

### Parasites of Concern

Cattle can be infected by roundworms (nematodes), tapeworms (cestodes) and flukes (trematodes). Protozoans such as coccidia are another type of internal parasite; however, the helminths (worms) will be the focus of this discussion.

Roundworms are considered the most economically devastating internal parasites of livestock in Arkansas. The medium or brown stomach worm and the *Cooperia* species are the most common roundworms. Although cattle can be infected with tapeworms, their effect on animal performance is minimal compared to the roundworms.

*Arkansas Is  
Our Campus*

Visit our web site at:  
<https://www.uaex.uada.edu>

Problems associated with flukes occur in some parts of Arkansas. Problems with flukes arise in conditions that promote snail populations – poorly drained pastures and stagnant pools of water (ponds, ditches, etc.) in the pasture area. Snails are needed in the fluke life cycle.

### Life Cycle of the Nematode

Figure 1 shows the life cycle of the nematode. In the host animal, adult nematodes produce eggs. The egg is expelled from the host with the feces contaminating the pasture. A first-stage larva hatches from the egg. The larva will molt two times before it becomes a third-stage larva. Once the larva is in its third stage, it is capable of migrating from dung pats and soil onto moist grass. Larvae can survive up to a year on pasture.

Infection occurs when the third-stage larva is consumed with the grass. The larva completes its life cycle in the gastrointestinal tract of its host. Once the adult stage is reached, copulation occurs and the life cycle starts over.

Unlike other nematodes, the medium stomach worm can spend part of its parasitic life cycle in hypobiosis, a condition similar to hibernation. Hypobiosis usually begins in the spring. The “hibernating” larvae do not emerge until summer.

### Life Cycle of the Fluke

The fluke’s life cycle requires two hosts – cattle and snails. The adult flukes are found in the bile ducts of cattle. The eggs are laid in the ducts and expelled with the feces. A larval stage hatches from the egg and infects the snail, where it reproduces asexually. Specific stages of the juvenile fluke leave

the snail and encyst on aquatic vegetation. Cattle eat the vegetation and become infected. The fluke migrates to the liver, infects the bile ducts and matures into an adult.

### Seasonal Parasite Pressure

The amount of parasite pressure in a pasture varies with season and management. Parasite burden peaks during the spring and is lowest during the hot, dry summer months (Figure 2). Cattle in drylot systems typically have fewer worms and less seasonal variation. Parasite pressure will be less under good management conditions as well. Good herd management includes a good nutrition and health program.





Figure 2. Seasonal Parasite Trends

### Parasite Infection Diagnosis

Parasite infection is diagnosed by either a fecal egg count or postmortem exam in the event of death. The postmortem examination is the most accurate method of determining parasitism. A postmortem exam shows the types of worms present and the damage to the animal.

Figure 1. Basic Life Cycle of Common Gastrointestinal Nematodes of Cattle



Internal parasite infection levels can be estimated in live cattle by counting the eggs shed with the feces. The eggs are quantified as eggs per gram (EPG) of feces by a trained individual. An EPG is only an estimate of parasite populations. Results from this test may be misleading. Variations in egg counts can arise from the ratio of immature to mature worm populations, worm species, degree of inhibition, consistency of the manure, etc. Realistically, a herd should be sampled on multiple occasions to determine worm prevalence.

## Pasture Practices for Reducing Parasitism

Pasture management and anthelmintics (dewormers) are two methods now used to control internal parasites. Pasture management practices may reduce the parasite burden in cattle; however, this method alone will not guarantee parasite eradication.

As discussed earlier, part of the nematode life cycle is on pasture. Pasture management methods designed to reduce third-stage larva populations include the following:

- Move more susceptible younger cattle to a safe pasture. Safe pastures include pastures that were not grazed during the last 12 months as well as small grain pastures developed from a prepared seedbed. When a pasture lies untilled and is plowed, contamination can drop quickly. Always deworm cattle prior to placement on a safe pasture; otherwise, the pasture can immediately become contaminated.
- Place less susceptible, mature cattle on the more contaminated pastures. Mature cows under a good nutrition program develop some acquired immunity to parasites and are affected less by their presence than young cattle and calves.
- Do not overgraze pastures. Animals on overgrazed pastures graze closer to the ground and pick up more larvae. Rotational grazing systems are unlikely to provide enough rest to paddocks to reduce possible contamination. Some studies have shown that rotational grazing can increase infection compared to continuous stocking. This is likely since rotational grazing allows higher stocking rates. However, rotational grazing combined with a strategic deworming program can still provide more production per acre than conventional grazing. If flukes are a problem, identify ways to increase pasture drainage and fence off problem areas such as ponds.

- Miscellaneous practices include dragging manure pats in dry weather and cutting the forage for haylage.

## Anthelmintic Control of Parasites

Anthelmintics provide an excellent tool for controlling parasites. Application of dewormers should not be aimed at treating infected cattle showing signs of parasitism. Instead, apply dewormers in a timely manner to reduce infection before symptoms of disease occur. Treatment should also be aimed at interrupting the life cycle of the parasite in an effort to minimize pasture contamination. Unfortunately, instead of implementing a deworming program, producers typically deworm their cattle when the herd is being worked for another purpose. The 2007-2008 NAHMS cow/calf health survey showed that 85 percent of beef cattle operations in the United States deworm their herd according to a regular schedule. Ten percent base treatment upon animal appearance. In the latter case, the herd has already suffered economic losses.

Many anthelmintic products are on the market. Most of the products are either avermectins/milbemycins (ivermectin, doramectin, eprinomectin and moxidectin) or benzimidazoles (oxfendazole, albendazole, fenbendazole). Avermectins/milbemycins provide an additional benefit of external parasite control plus internal parasite control, as well as persistent protection for days to weeks after treatment. When selecting a dewormer, the following should be kept in mind:

- Type of animal being treated (calf vs. cow, beef vs. dairy)
- Product efficacy
- Ease of application
- Broad spectrum of control (immature, mature, inhibited)
- Cost effectiveness
- Slaughter/milk withdrawal time
- Personal safety

Table 1 is a summary of the common products currently available. Table 2 summarizes the effectiveness of the different active ingredients. Remember, the production situation on every farm is different. Evaluate the pros and cons of all products before choosing a dewormer. If possible, conduct a fecal egg count before and after deworming to evaluate the effectiveness of the program.

**Table 1. Commonly Used Deworming Products**

Type	Trade Name	Product* Active Ingredient	Warnings and Withdrawals**	
			Dairy & Milk	Beef & Slaughter
<b>Block</b>	Safe-Guard Cattle Block	Fenbendazole	Note 1	11d
<b>Drench</b>	Prohibit	Levamisole	Note 1	48h
	Synanthic 9.06%	Oxfendazole	Note 1	7d
	Synanthic 22.5%	Oxfendazole	Note 1	7d
	Panacur	Fenbendazole	0	8d
	Safe-Guard	Fenbendazole	0	8d
	Valbazen	Albendazole	Note 1, 2	27d (Note 2)
<b>Feed Additives</b>	Safe-Guard	Fenbendazole	0	13d
	Rumatel	Morantel Tartrate	0	14d
<b>Injectable</b>	Levamisol	Levamisole	Note 1	7d
	Ivermectin Containing	Ivermectin	Note 1	35d
	Ivomec Plus, Noromectin Plus	Ivermectin/Clorsulon	Note 1	49d
	Dectomax	Doramectin	Note 3	35d
	Cydectin	Moxidectin	Note 1	21d
<b>Paste</b>	Panacur	Fenbendazole	0	8d
	Safe-Guard	Fenbendazole	0	8d
<b>Pour-on</b>	Ivermectin Containing Pour-on	Ivermectin	Note 1	48d
	Eprinex	Eprinomectin	0	0
	Dectomax Pour-on	Doramectin	Note 3	45d
	Cydectin Pour-on	Moxidectin	0	0
<b>Mineral</b>	SafeGuard Mineral dewormer	Fenbendazole	0	13d

\*Local feed dealerships may independently market feed mixes and blocks containing additive products.

Note 1: Not to be used on dairy cattle of breeding age.

Note 2: Do not administer during the first 45 days of pregnancy or for 45 days after bull removal.

Note 3: Safe in dairy heifers up to 20 months of age.

**\*\*Withdrawals are subject to change; always read the label before purchasing an animal health product.**

**Table 2. Major Parasiticides and Their Efficacies**

Anthelmintic	Parasite							
	Roundworm		Liver Fluke	Tape- worm	Mites	Lice	Warbles	Persistence
	Inhibited	Active						
Levamisole	None	Fair	None	None	None	None	None	None
Oxfendazole	Fair	Good	None	Some	None	None	None	None
Fenbendazole	Fair	Good	None	Some	None	None	None	None
Albendazole	Fair	Good	Some	Some	None	None	None	None
Avermectins	Good	Good	None	None	Good	Good	Excellent	Good
Milbemycins	Good	Good	None	None	Good	Good	Excellent	Good
Clorsulon	None	None	Good	None	None	None	None	None

Always read and follow the instructions on any animal health product. Application rates were developed through extensive research to identify the best and safest level of control. Undertreatment of animals can reduce the level of control and may increase the chances of parasites developing resistance to dewormers. Applying dewormers above recommended levels increases withdrawal times, and the compounds in dewormers can become toxic to animals when applied at extremely high rates. Overdosing is costly. Over application usually provides the same amount of control as recommended levels but at a higher cost.

As with any chemical used to control parasites, the potential for parasites to develop resistance exists. No reports of extensive parasite resistance to currently used bovine anthelmintics exist in the U.S.; however, countries with livestock numbers comparable to the U.S. have reported occurrences of resistance in areas where dewormers were used extensively.

## Deworming the Beef Herd

Since cows, bulls and young stock are affected differently by internal parasites, corresponding treatment programs should also differ. Table 3 gives a recommended guideline for deworming beef cattle.

Mature cows should be treated at least one time per year. The best time to treat the mature cow is near calving. The mature cow's susceptibility to parasite detriment increases during this time due to stress of production and a suppressed immune system. In situations where parasite levels may be high, such as overstocked pastures, treating twice a year may be necessary. In other situations, parasite levels may be low enough that mature cows do not need any treatment. These conditions can only be determined by treatment followed by critical observation. Bulls, unlike cows, tend to be more susceptible to parasites and should be treated twice a year, spring and fall. Since mature cows and bulls have some degree of acquired immunity to internal parasites, the older benzimidazole-type products should do a sufficient job of controlling parasites in these animals.

Treatment of calves should begin when they reach three to four months of age and again at weaning if they are kept as replacements or stockers. Yearlings can be treated on a seasonal basis, spring and fall, until they are mature cows (a mature cow is generally recognized as an animal pregnant with her second calf). If calves are backgrounded in a drylot, one initial treatment should be sufficient.

Visit with a local veterinarian about setting up a herd health program that includes a strategic deworming program.

**Table 3. Deworming Program for Beef Cattle**

Animal	Time of Treatment
Mature cows	Near calving
Bulls	Spring and fall
Calves	3 to 4 months of age
Replacements and stockers	<ul style="list-style-type: none"> <li>• Weaning/purchase and at spring/fall (minimum)</li> <li>• Weaning/purchase and every 3 to 4 months until yearlings</li> <li>• Weaning/purchase and placed on safe pasture</li> </ul>
Yearlings	Spring and fall until mature

## Deworming the Dairy Herd

Deworm dairy heifers and bulls on pasture according to the recommendations for beef calves and replacements. Initial deworming of calves should occur three to four weeks after turnout onto pasture in the spring.

If calves or replacements are raised in total confinement or drylot, take egg counts to establish the level of parasitism and design a deworming program accordingly. Young calves weighing less than 400 pounds will need to be dewormed more frequently than 800-pound yearlings, as younger calves are more susceptible to worms than older animals.

The first step in designing a deworming program for lactating dairy cows is to determine the level of parasitism. Research data shows variable responses to deworming for mature dairy cattle. Deworming at early lactation increases milk yields more consistently than any other time of treatment.

Parasite contamination potential in dairy cattle can be grouped into four categories: high, moderate, low and extremely low. High contamination potential exists in dairy cattle on pasture during lactation and when rotational grazing is practiced; whereas, an extremely low potential exists for cattle confined to a drylot. Table 4 is a guideline for treating cattle based on the different levels of exposure.

**Table 4. Deworming Program for Dairy Cows**

Level of Contamination	Time of Treatment	Recommended Product Type
High to moderate	Freshening	Avermectin or Benzimidazole
Low	Freshening	Avermectin or Benzimidazole
Extremely low	No treatment	

Since dairy cows are most likely to show a positive response to deworming during early lactation, the first choice of a deworming program is to treat at freshening. If at all possible, a producer should treat some cattle and monitor the production of all cattle. This would serve to quantify the importance of worms in the operation.

In high production (over 18,000 pounds of milk/cow/year and seasonal calving in the fall), another option is to deworm all cows at one time and then again six to eight weeks later. Another option for

herds with high and moderate levels of contamination is to deworm in the fall and again in the spring. This latter option may not be economical since cows with low parasite loads and cows in mid- to late lactation may not respond with an increase in milk production after treatment. Use only products approved for use in lactating dairy cattle, and consider milk withdrawal times when treating.

Visit with a local veterinarian about setting up a herd health program which includes a strategic deworming program.

The information given herein is for educational purposes only. Reference to commercial products or trade names is made with the understanding that no discrimination is intended and no endorsement by the University of Arkansas Division of Agriculture is implied.

Printed by University of Arkansas Cooperative Extension Service Printing Services.

**DR. SHANE GADBERRY** is associate professor - animal science, Little Rock, and **JEREMY POWELL, DVM**, is associate professor - veterinarian, Fayetteville. They are employees of the Department of Animal Science, University of Arkansas Division of Agriculture.

Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Director, Cooperative Extension Service, University of Arkansas. The Arkansas Cooperative Extension Service offers its programs to all eligible persons regardless of race, color, national origin, religion, gender, age, disability, marital or veteran status, or any other legally protected status, and is an Affirmative Action/Equal Opportunity Employer.