

Home Gardening Series
Brussels Sprouts

Craig R. Andersen
Associate Professor

Environment

Light – sunny
Soil – well-drained, loam
Fertility – rich
pH – 5.8 to 6.8
Temperature – cool
Moisture – keep moist



Culture

Planting – **spring:** transplant early or
fall: sow seeds early to
midsummer or transplant
late summer
Spacing – 12-18 x 24-30 inches
Hardiness – hardy biennial
Fertilizer – heavy feeder

Brussels Sprouts – *Brassica oleracea* gemmifera

Brussels sprouts are a variety of cabbage. Their existence was first recorded in 1587, and they were apparently developed in the 15th century in the northern part of Europe that is now Belgium. Whether they originated in Brussels or not, they were cultivated there for centuries and rapidly spread to other countries around the world. In England they are a popular vegetable served with holiday meals. Brussels sprouts first became popular in

Europe after World War I, when consumption of vegetables increased considerably, but they are still among the minor vegetables in the United States.

Brussels sprouts (named after Brussels, Belgium, where the vegetable was first popular) are hardy, slow-growing, long-season vegetables that belong to the cabbage family. They can be grown with success in Arkansas if planted very early in the spring or in late summer for fall production.

Cultural Practices

Planting Time

These are cool-season plants that thrive during the long periods of cool weather we have in the fall and early winter. Use transplants in late winter for a late spring harvest. Fall planting is generally the best. Sow seed in a protected location in the garden four

*Arkansas Is
Our Campus*

Visit our web site at:
<https://www.uaex.uada.edu>

Cultivars

Cultivar	Days to Maturity	Plants/ 100 Ft of Row	Disease Resistance or Tolerance	Remarks
Bubbles	90	66		Hybrid, 2-foot tall plant, best for fall planting but holds well in spring.
Jade Cross Hybrid	95	50	botrytis	Uniform maturity, good yields.
Long Island Improved	95	50		Large plant, good yields.

or five weeks before transplanting. Transplant the seedlings to the permanent garden location. For fall harvest, transplant in early to midsummer, about the same time you would plant late cabbage.

When transplanting, use 8 ounces per plant of a starter fertilizer solution (1 tablespoon of 20-20-20 soluble fertilizer per gallon of water).

Spacing and Depth of Planting

Space plants 24 inches apart in the row. Cover the seed 1/4 to 1/2 inch deep. Transplant seedlings when they are about 3 inches tall.

Care

Brussels sprouts are grown much like related cole crops – cabbage and broccoli. Apply one side-dress application of nitrogen fertilizer when the plants are 12 inches tall and then every three to four weeks. Water to keep the plants growing vigorously during the heat of summer. Without ample soil moisture, the crop will fail. Cultivate shallowly around the plants to prevent root damage. The sprouts form in the axils of the leaves (the space between the base of the leaf and the stem above it).

Side-dress the plants with a nitrogen fertilizer, 10-20-10 or 13-13-13, when they are half grown. On sandy soils or after periods of excessive rain, fertilize every three weeks. Use shallow cultivation and mulches to control weeds.

Commercial gardeners top the plants (removing the apical meristem) to promote the growth of the sprouts once the plants have reached a height of 24 to 36 inches. The spring crop should be shorter to hasten development before it gets too hot. Gardeners will remove the lower leaves to expose the sprouts. They believe the sprouts develop better if the

lowermost six to eight leaves are removed from the side of the stalk as the sprout develops. Two or three additional leaves can be removed each week, but several leaves should be left intact on top. Removing the lower leaves will accelerate harvest, but this is not necessary in the home garden.

Common Problems

Diseases

Yellows or fusarium wilt is a common disease. Yellowing and browning of lower leaves is the first sign of the disease. The plant will be stunted before wilting occurs. Grow varieties resistant to yellows.

Blackleg and blackrot are two bacterial diseases that cause severe losses. The plant may be stunted, turn yellow and die; the taproot often rots away. Blackleg is named for the black cankers on the stem. Blackrot can be recognized by large, V-shaped, yellow-to-brown areas in the leaves, starting at the leaf edge. The veins turn black. Soft rot usually follows blackrot infection. Control of blackleg and blackrot is essentially the same. Both diseases are spread by seeds, transplants and insects. Buy seeds that have been treated with hot water to kill the disease organism. Do not buy transplants that are wilted, are an unhealthy shade of green or have black spots on the stems or leaves. When diseased plants are found in the garden, collect the entire plant and dispose of it. Do not put diseased plants in the compost pile. Avoid cultural practices (crowding, overwatering, planting in poorly drained soil and inadequate insect control) that support blackrot and blackleg disease organisms.

Insects

Aphids – Apply a suggested insecticide before Brussels sprouts begin to head.

Flea Beetles – Apply a suggested insecticide.

Cabbage Worms – Three species of cabbage worms (imported cabbage worm, cabbage looper and diamondback worm) commonly attack the leaves and heads of Brussels sprouts and related cole crops in Arkansas. Imported cabbage worms are velvety green caterpillars. The cabbage looper crawls by doubling up (to form a loop) and then moves the front of its body forward. The moth is brown and is most active at night. Diamondback worms are small, pale green caterpillars that are pointed on both ends. The moth is gray with diamond-shaped markings when the wings are closed.

Larval or worm stages of these insects cause damage by eating holes in leaves. The adult moths or butterflies lay their eggs on the leaves but otherwise do not cause damage to the plant. The worms are not easy to see because they are very small and blend in with the leaves. Cabbage worms are quite destructive and will ruin a crop if not controlled. They are more destructive in fall gardens than in spring gardens. Loopers, a severe pest of Brussels sprouts, are easy to control using the biological insecticide *Bacillus thuringiensis* (B.t.). This material gives excellent control of worms and can be used with complete safety around the home. It is sold under many trade names such as Thuricide, Dipel and B.t. Use one to two drops of a liquid detergent per gallon of spray mixed to ensure complete wetting of the waxy leaf surface. Protect your plants with suggested biological or chemical insecticides from the time they are transplanted until harvest.

Harvesting

The small sprouts or buds form heads 1 to 2 inches in diameter. They may be picked (or cut) off the stem when they are firm and about 1 inch in size. The lower sprouts mature first. The lowermost leaves, if they have not already been removed, should be removed when these are harvested. Harvest the sprouts before the leaves turn yellow.

approximate yields (per 10 feet of row) – 10 to 18 pounds

amount to raise per person – 10 pounds

storage – very cold (32 degrees F), moist (95 percent relative humidity) conditions, 4 to 6 weeks

preservation – freeze



Frequently Asked Questions

Q. Why do my sprouts remain loose tufts of leaves instead of developing into firm heads?

A. When the sprouts develop in hot weather (this may be spring or fall), they often do not form compact heads. That is why they are a better fall and winter crop. Use transplants for early planting and maintain ample soil moisture.

Q. When do I plant Brussels sprouts for maximum production?

A. Brussels sprouts are sensitive to temperature. In general, Brussels sprouts produce best when daytime temperatures average about 65 degrees F or less. Brussels sprouts grow best when planted in mid to late summer for late fall or early winter harvesting.

Q. Should you pinch or cut the top out of Brussels sprout plants to make them produce more?

A. In the spring, pinching or removing the growing point of the plant will hasten the development of the sprouts resulting in earlier harvest but will reduce the yield by about one-third.

Printed by University of Arkansas Cooperative Extension Service Printing Services.

DR. CRAIG R. ANDERSEN is associate professor, Department of Horticulture, University of Arkansas Division of Agriculture, Fayetteville.

Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Director, Cooperative Extension Service, University of Arkansas. The Arkansas Cooperative Extension Service offers its programs to all eligible persons regardless of race, color, national origin, religion, gender, age, disability, marital or veteran status, or any other legally protected status, and is an Affirmative Action/Equal Opportunity Employer.

FSA6005-PD-5-09RV