

Crapemyrtle Bark Scale: A New Insect Pest

James Robbins
Professor and Extension
Horticulture Specialist -
Ornamentals

John Hopkins
Associate Professor
and Extension
Entomologist - Urban

Mike Merchant
Professor and Extension
Urban Entomologist
Texas A&M AgriLife
Extension Service

Mengmeng Gu
Assistant Professor and
Extension Ornamental
Horticulturist
Texas A&M AgriLife
Extension Service

*Arkansas Is
Our Campus*

Visit our web site at:
<https://www.uaex.uada.edu>

History

A relatively new insect is appearing on crapemyrtles (*Lagerstroemia*) across the Southeast, including Arkansas (Figure 1). The insect was first noticed in a north Dallas, Texas, suburb in 2004. At that time, entomologists suggested it was morphologically identical to azalea scale (*Eriococcus azalea*), but the scientists noted that molecular investigation might eventually identify it as *E. lagerstroemiae*, known to be a pest on crapemyrtle and pomegranate (*Punica*) in Asia. Although the exact taxonomy is still not known, the insect is most commonly referred to as crapemyrtle bark scale (CMBS).

Since the initial sighting in 2004, the insect has been spreading at an alarming rate across the Southeast. The insect was first noted in McKinney, Texas (self-designated as America's "crapemyrtle city"), in 2005 and had spread throughout most of the Dallas-Fort Worth area by 2010. The scale was reported in Ardmore, Okla., and Shreveport, La., in 2012 and Houma, La. (60 miles southwest of New Orleans) in 2013. In October 2013, the insect was confirmed in Germantown, Tenn. (Memphis area), and in Little Rock, Ark., in January 2014.

Heightened concern about this new pest is based on the speed at which it is spreading and the common



Figure 1. Crapemyrtle bark scale (CMBS)

use of crapemyrtles in landscapes across a large part of the U.S. To date there has been no research to evaluate whether there is a range in susceptibility to this insect across the hundreds of crapemyrtle cultivars; however, anecdotal observations from McKinney, Texas, suggest the scale may be worse on hybrids (*fauriei* × *indica*).

Insect Description

Crapemyrtle bark scale is easy to identify since, in the U.S., it is the first and only known bark scale to occur on crapemyrtles. The adult



Figure 2. Adult female CMBS on crapemyrtle bark



Figure 3. Heavy infestation of CMBS on most recent flush of growth



Figure 4. Sooty mold from heavy CMBS infestation

females appear as white or gray felt-like encrustations (Figure 2) on small twigs to large trunks, often appearing near pruning wounds or in branch crotches on older wood. On the most current flush of growth and under heavy infestation, distribution may be more uniform (Figure 3). Up close, CMBS is white to gray in color and approximately 2 mm in length. Careful examination may reveal dozens of pink eggs or crawlers under some of the larger white scale covers. Most gardeners will be alerted to CMBS by black sooty mold which appears on the bark (Figure 4). The presence of sooty mold may confuse the diagnosis since that is also commonly associated with a significant aphid problem. This felt scale is not classified as either an armored or soft scale.

Life Cycle

As female nymphs mature, they secrete white threads (Figure 5) that become felted or matted into a thick, whitish to grayish scale covering over the entire body (Figure 6). Adult females under this covering are wingless and sessile (attached and incapable of moving). It is thought that for this species of scale, eggs are laid under the covering (probably late April to mid-May in Arkansas) and the female then dies. When the eggs hatch into first instar nymphs, these nymphs have legs and antenna and are mobile, thus the term *crawlers* (Figure 7). These crawlers emerge from under the “mother scale” and disperse over a short period (about one to two days). We suspect that this emergence occurs beginning mid-May to early June in Arkansas; however, crawlers

were observed on a warm day in January 2014. After the first molt, nymphs lose their legs and antenna and become sessile (Figure 5). During the last instar, males are quiescent (pupal type stage) and develop external wings. Upon emergence, males find a sessile female and mate, completing the life cycle. At present in the Southern U.S., the number of generations completed in a year for this species is unknown, but it is suspected that CMBS may complete at least two generations in Arkansas. It is possible that adult females and eggs overwinter, but crawlers and later stage nymphs have been observed overwintering in Arkansas under loose bark and in cracks and crevices (Figure 8).



Figure 5. CMBS nymphs exuding white threads that ultimately form the felt cover



Figure 6. Adult female CMBS



Figure 8. Settled CMBS nymphs under loose bark



Figure 7. CMBS crawlers on pruning cuts (L) and close-up of crawlers (R)

Control

Based on our limited experience with this pest, it does not appear that CMBS will be easy to control, though soil-applied neonicotinoids do provide significant suppression. Our current best suggestions for control of this insect include:

- For heavily infested plants, wash the trunk and reachable limbs with a soft brush and mild solution of dishwashing soap. This will remove many of the female scales and egg masses and make insecticide control more effective. Also, washing will remove much of the black mold that builds up on the bark on infested trees.
- Horticultural oil has not yet been shown to be effective against this insect; however, a winter application of **dormant oil** to the bark and crotches of the plants where scales shelter may be beneficial. Be sure to use sufficient volume to allow for penetration behind loose bark and into cracks and crevices. Winter is an especially good time to treat for scales because a higher (winter) application rate can be used without damaging the plant. Thorough coverage of the tree is especially important when treating with oil.
- Application of systemic insecticides as a drench applied to the root zone has shown the most promise in tests to date. Imidacloprid (Merit® or Bayer Advanced™ Garden Tree and Shrub Insect Control), thiomethoxam (Meridian®) and dinotefuran (Greenlight Tree and Shrub Insect Control with Safari) have shown best control when applied between May and July. When drenching the soil with a systemic insecticide, allow several

weeks for the product to be distributed throughout the plant. Additionally, acetamiprid and clothianidin, also neonicotinoids, have demonstrated good control.

- Certain insect growth regulators (IGRs) are recommended for scale control in woody ornamentals (MP144, *Insecticide Recommendations for Arkansas*) but have not yet been evaluated on CMBS.
- Lady beetles should be preserved, as the twice-stabbed lady beetle is an efficient predator of this scale (Figure 9).



Figure 9. Pupa of lady beetle, predator of CMBS

If you notice symptoms of scale insect infestation on your crapemyrtles in Arkansas, please contact the office of your local Cooperative Extension Service or the Arkansas State Plant Board.

References

- Merchant, M. Crape myrtle scale: New pest of U.S.? <http://citybugs.tamu.edu/2010/05/10/scale/>. Accessed on Jan. 10, 2014.
- Merchant, M. Update on crape myrtle scale. <http://crapemyrtletrails.org/pest-control/>. Accessed on Jan. 10, 2014.
- Morgan, J. Invasive species troubles crape myrtles. http://www.lsuagcenter.com/news_archive/2013/october/headline_news/Invasive-species-troubles-crape-myrtles.htm. Accessed on Jan. 10, 2014.

All chemical information provided is given with the understanding that no endorsement of named products is intended nor is criticism implied of similar products that are not mentioned. Individuals who use pesticides are responsible for ensuring that the intended use complies with current regulations and conforms to the product label. Before purchasing or using any pesticide, always read and carefully follow the label directions.

DR. JAMES ROBBINS is professor and Extension horticulture specialist and **DR. JOHN HOPKINS** is associate professor and Extension entomologist - urban, University of Arkansas Division of Agriculture, Little Rock. **DR. MIKE MERCHANT** is professor and Extension urban entomologist and **DR. MENG MENG GU** is assistant professor and Extension ornamental horticulturist, Texas A&M AgriLife Extension Service, College Station. FSA7086-PD-1-14N

Pursuant to 7 CFR § 15.3, the University of Arkansas System Division of Agriculture offers all its Extension and Research programs and services (including employment) without regard to race, color, sex, national origin, religion, age, disability, marital or veteran status, genetic information, sexual preference, pregnancy or any other legally protected status, and is an equal opportunity institution.