

**Crop Replant and Rotation Guide for Soybean Herbicides\***

| Herbicide                                   | Replant/Crop Rotation                                    | Time Interval   | Precautions  |
|---|--|---|--|
| Anthem Maxx                                 | S,C<br>CT,P,SF,W<br>AL,R<br>SG                           | I<br>4 months<br>10 months<br>11 months                                       | See label for rotational intervals based on rates applied.   |
| Assure II                                   | S,C<br>All   | I<br>120 days   |  |
| Authority MTZ                               | S<br>B,W,SG<br>R<br>AL,CT,GS,P,SF                        | I<br>4 months<br>10 months<br>12 months                                       | See label for pH restrictions. High pH = longer rotations.   |
| Authority Edge/<br>Authority Supreme        | S, SF<br>C,P,W<br>GS,R<br>B<br>AL,CT,O                   | I<br>4 months<br>10 months<br>11 months<br>12 months                          | See label for pH, rate restrictions. High pH rates have longer rotational intervals.   |
| Authority XL                                | B,R,Y,W<br>AL,CL,CT,<br>FC,GS,O,<br>P,R,SC,SF<br>All     | 4 months<br>18 months<br><br>36 months  | See label for pH, rate restrictions.   |
| Basagran                                    | All  | I   |  |
| Broadaxe XC or<br>Authority Elite           | S,SU,DSP<br>P<br>W<br>FC,R,GS<br>O,CT*<br>SC,PC          | I<br>4 months<br>4.5 months<br>10 months<br>12 months<br>18 months            |  |
| Butyrac/Butoxone<br>(2,4-DB)                |  |   | No restrictions.   |
| Cadet                                       | All  | I   |  |
| Canopy                                      | S<br>W,B,RG<br>C<br>AL,CT,GS,R<br>CL<br>P<br>All         | I<br>4 months<br>9 months<br>10 months<br>12 months<br>8 months<br>18 months† | Use only on soils with pH of 7.0 or less.<br>† Successful field bioassay must be completed prior to planting. Wait 15 months before trying a bioassay.   |
| Canopy (pH > 7)<br>Do not use<br>> 3.5 oz/A | S<br>W,B,RG<br>C<br>AL,CT,GS<br>CL<br>P<br>R, all others | I<br>4 months<br>9 months<br>10 months<br>12 months<br>8 months<br>18 months  | Use only in soils with pH 7.5 or less.<br>† Successful field bioassay must be completed prior to planting. Wait 15 months before trying a bioassay.  |
| Canopy EX                                   | W<br>C,GS,R,AL,CL<br>CT<br>SF,CA                         | 3 months<br>9 months<br>10 months<br>18 months                                | See label for pH, rate and geographical restrictions. High pH soils have longer rotational intervals.  |
| Classic                                     | S†<br>SG,RG<br>P<br>C,CT<br>CL,GS,R<br>All               | I<br>3 months<br>6 months<br>8 months<br>9 months<br>18 months††              | If applied after Aug. 1, extend recrop interval for C, CT, R and GS 2 months.<br>† Recrop intervals may also be extended if Classic is applied after application of Canopy. See labels for information.<br>†† Successful field bioassay must be completed prior to planting. Wait 9 months before trying a bioassay. |

| Herbicide                  | Replant/Crop Rotation   | Time Interval   | Precautions   |
|----------------------------|---|---|---|
| Cobra                      |   |   | No restrictions.  |
| Dual Magnum or<br>Sequence | S,C,CT,GS†<br>SG,W<br>Rice<br>All                                       | I<br>4.5 months<br>Next spring<br>18 months                           | † Use Concep-treated grain sorghum seed.  |
| Enlist One                 | S, C<br>CT<br>All   | 7-14 days<br>1 month<br>1 month                                       | Rotation intervals based on rate – see label. Enlist corn, cotton, and soybean can be planted immediately.  |
| Enlist Duo                 | S, C, CT<br>All   | 1 month<br>1 month  | Enlist corn, cotton, and soybean can be planted immediately.  |
| Engenia                    | C<br>GS, S, SG<br>All   | I<br>14-28 days<br>120 days   | Rotation intervals based on rate – see label. Dicamba-tolerant soybean and cotton can be planted immediately.   |
| Enlite                     | W<br>CT,C,R,SF<br>AL,CL   | 4 months<br>9 months<br>12 months                                     |   |
| Envive                     | W<br>CT,C,GS,R,AL,CL<br>SF,CA   | 4 months<br>10 months<br>18 months                                    | See label for pH and geographical restrictions. High pH soils have longer rotational intervals.   |
| FirstRate                  | W<br>C,CT,P,R,GS  | 3 months<br>9 months  | All others – 30 months plus bioassay.   |
| Flexstar GT 3.5            | CT,S<br>W<br>C,P,R,GS<br>Other  | 0 months<br>4 months<br>10 months<br>18 months                        | See Reflex.   |
| Fusilade DX or<br>Fusion   | S, CT<br>All  | I<br>2 months   |   |
| Glufosinate                | C,S,CT,R,CA<br>W,B,O<br>GS  | I<br>70 days<br>180 days  | No restrictions.  |
| Glyphosate                 | All   | I   |   |
| Intimidator                | S<br>B, W<br>C, R<br>CT, RY<br>GS<br>All                                | I<br>4.5-8 months<br>10 months<br>12 months<br>18 months<br>18 months |   |
| Metribuzin                 | S†<br>AL,C,FG,W,B<br>CT,R<br>All (except root<br>crops)††<br>Root crops | I<br>4 months<br>8 months<br>12 months<br>18 months                   | † Waiting period for replanting soybeans depends on the rate of metribuzin used. See specific label for more information. Add 2 months to time intervals if pH of soil is above 7.5.<br>†† Cover crops may be planted any time, but stand reductions may occur. |

\*This table applies to the major field and forage crops. Refer to the herbicide labels for the latest recrop and rotation information for horticultural crops. These are written as best we could interpret the labels. We regret any omissions or errors. Always refer to product labels before using a pesticide or replanting into treated fields.

NOTE: See page 44 for key to abbreviations.

*(continued on page 44)*

**Crop Replant and Rotation Guide for Soybean Herbicides\* [cont.]**

| Herbicide             | Replant/Crop Rotation   | Time Interval  | Precautions  |
|-----------------------|---|--|--|
| Outlook               | C,S<br>SG<br>All  | I<br>4 months<br>FY  | Do not retreat field with second application.  |
| Paraquat              |   |  | No restrictions.   |
| Poast/Poast Plus      |   |  | See label.   |
| Prefix                | S<br>CT<br>AL,W,O,R,RY,B<br>GS,R,C                                  | I<br>1 month<br>4.5 months<br>10 months  | Use Concep-treated grain sorghum seed. Do not use Reflex or Flexstar in-season.  |
| Prowl                 | S,CT<br>W,B<br>All  | I<br>4 months†<br>FY   | Do not rework soil deeper than treated zone.<br>† Cannot replant using no-tillage practices.   |
| Pursuit               | S<br>B,W<br>C<br>Rice<br>All  | I†<br>4 months<br>9.5 months<br>18 months<br>18 months   | † Do not rework soil deeper than 2 inches. Do not apply Classic, Canopy, Lorox Plus, Scepter, Squadron or Tri-Scepter the same year as Pursuit or injury to following crops may occur. |
| Python                | AL,P,B,W,RY,O<br>R<br>C<br>GS<br>CT, Sunflower                      | 4 months<br>6 months<br>1 month<br>12 months<br>18 months  | Requires successful bioassay.  |
| Reflex or Flexstar    | S<br>SG, W<br>C,CT,R,P,GS<br>All                                    | I<br>4 months<br>10 months<br>18 months  |  |
| Resource              | SG<br>S,C<br>All  | I<br>120 days<br>30 days   |  |
| Select Max            |   |  | See label.   |
| Sharpen               | C,GS,SG,S,R<br>CT<br>SF, others                                     | I<br>1.5 months<br>4 months  | † Rotation intervals based on rate – see label.  |
| Sonic/Authority First | W<br>C,R<br>CT,AL,B,O, RY,GS  | 4 months<br>10 months<br>12 months   | See label for pH restrictions.   |
| Storm                 | All   | Fall   |  |
| Surveil               | S<br>W<br>C,P,R,GS<br>SF  | I<br>3 months<br>9 months<br>30 days   |  |
| Tavium                | S<br>CT<br>C<br>B, O, RY, W<br>AL, DSP, GS, P, SF<br>CL<br>R<br>All | 28 days<br>42 days<br>4 months<br>4.5 months<br>6 months<br>9 months<br>Next spring<br>12 months | Dicamba-tolerant soybean and cotton can be planted immediately.  |
| Treflan/others        | S,CT<br>W,B<br>All  | I<br>Fall<br>FY  |  |

| Herbicide         | Replant/Crop Rotation                              | Time Interval   | Precautions  |
|-------------------|--|---|--|
| Treflan (2X rate) | S,CT<br>R<br>All                                   | I<br>FY<br>2 years  |  |
| Trivence          | S<br>W<br>C<br>R<br>CT<br>All                      | I<br>4 months<br>10 months<br>12 months<br>18 months<br>18 months               | Rotational intervals based on pH and rates – see label.  |
| Typhoon           | W<br>C,CT,R<br>GS and all others                   | 4 months<br>10 months<br>18 months  |  |
| Ultra Blazer      | S<br>All   | I<br>Fall   |  |
| Valor             | S,P<br>W,C,R,GS<br>CT<br>B<br>AL,O,CL,CA           | I<br>30 days<br>21 days<br>4 months<br>8 months                                 | Pre-emerge treatments may injure soybeans.   |
| Verdict           | C,S,GS<br>CT,R<br>W                                | I<br>FY<br>4 months   | † Rotation intervals based on rate – see label.  |
| Warrant           | CT,C,GS,S<br>W<br>R, others                        | I<br>4 months<br>FY   | Do not use more than 3.0 lb/A/year.  |
| Xtendimax         | C, S<br>GS<br>CT<br>SG<br>R<br>All                 | 14-28 days<br>15 days<br>21 days<br>22-45 days<br>120 days<br>120 days          | Rotation intervals based on rate – see label. Dicamba-tolerant soybean and cotton can be planted immediately.  |
| Zidua             | CT,C,S<br>W<br>GS<br>R, others                     | I<br>1 month<br>6 months<br>FY  |  |
| Zidua Pro         | S<br>P, W<br>C<br>AL<br>B, RY<br>CT, GS, SF<br>All | I<br>4 months<br>8.5 months<br>10 months<br>11 months<br>18 months<br>40 months | Following 40 months, before planting any crop not listed on the label, a field bioassay must be completed. See label for Clearfield crop rotation intervals. |

**KEY**

**Crop**

All = All crops not specified  
 AL = Alfalfa  
 B = Barley  
 C = Corn  
 CA = Canola  
 CL = Clovers  
 CT = Cotton  
 DSP = Dry Shelled Peas  
 FC = Field Corn

FG = Forage Grasses  
 FL = Forage Legumes  
 GS = Grain Sorghum  
 O = Oats  
 PC = Peanuts  
 P = Popcorn  
 R = Rice  
 RG = Ryegrass  
 RY = Rye

**Timing**

I = Immediately  
 FY = Following year (usually spring or following fall - 11 to 16 months)

\*This table applies to the major field and forage crops. Refer to the herbicide labels for the latest recrop and rotation information for horticultural crops. These are written as best we could interpret the labels. We regret any omissions or errors. Always refer to product labels before using a pesticide or replanting into treated fields.