



## Arkansas Plant Health Clinic Newsletter

Follow us on social media



### Corn

#### Corn Crazy Top

Crazy Top of corn, causal agent *Sclerophthora macrospora*, is widespread in the United States. This disease is one of the several downy mildews that attack corn and sorghum. Symptoms depend on the time of infection and degree of colonization by the pathogen. Excessive tillering (six to ten tillers per plant), rolling, and twisting of the upper leaves, and leafy proliferation of the tassel are common symptoms. Leaves may be narrow, straplike, leathery, chlorotic, and stunted. The oospores are 45-75µm in diameter, hyaline to yellow, globose, with granular contents. Sporangia are lemon shaped 30-65 x 60-100, attached to short, simple sporangiophores emerging from stomata. Crazy Top is a problem when soils have been flooded shortly after planting or before plants are in the four to five leaf stage. Water trapped in the whorl of small plants can also lead to infection. Soil or leaf saturation for 24-48 hours is enough for infection to occur. There are no chemical controls for Crazy Top. Good soil drainage is the only preventative.

#### Corn Crazy Top-*Sclerophthora macrospora*



Photo by Sherrie Smith, University of Arkansas Cooperative Extension

#### Corn Crazy Top-*Sclerophthora macrospora*



Photo by Sherrie Smith, University of Arkansas Cooperative Extension

The University of Arkansas System Division of Agriculture offers all its Extension and Research programs to all eligible persons without regard to race, color, sex, gender identity, sexual orientation, national origin, religion, age, disability, marital or veteran status, genetic information, or any other legally protected status, and is an Affirmative Action/Equal Opportunity Employer.



## **Corn Crazy Top oospores- *Sclerophthora macrospora***



Photo by Sherrie Smith, University of Arkansas  
Cooperative Extension

## **Corn with Branched Ears**

Branched ears develop when pollination of the main ear is poor. This is usually attributable to weather factors at pollen shed.

## **Corn with Branched Ears-Abiotic**



Photo by Sherrie Smith, University of Arkansas  
Cooperative Extension



## Cotton

Bacterial Blight is a major disease of cotton in most of the cotton production areas of the world. The causal agent of Bacterial Blight is *Xanthomonas campestris* pv. *malvacearum*, an aerobic, Gram-negative rod-shaped bacterium. The disease can attack stems, leaves, petioles, and pods. The first symptoms on seedlings are small, water-soaked circular or irregular spots that turn brown. Spots on the hypocotyls are black, elongate lesions that often girdle the stem and cause seedling death. Leaf lesions first appear as minute water-soaked spots, becoming larger, angular, and black. Bacterial oozing may be observed during wet, humid conditions. Often, vascular infections follow the main ribs of the leaf, causing a black discoloration along the veins. Cotton Bacterial Blight is known as Black Arm when the disease extends from infected leaves into the petiole and stem. This can cause girdling of the stem, followed by stem death and stem breakage. Symptoms on the bolls begin as round water-soaked raised lesions that become irregular in shape and sunken and turn brown to black. Secondary infections can cause severe boll rots. Crop residue should be plowed under to speed decomposition and death of the bacterium. The use of acid-delinted and fungicide-treated seed, good sanitation, and the use of resistant cultivars lessen crop losses from Bacterial Blight.

## Cotton Bacterial Blight- *Xanthomonas campestris* pv. *malvacearum*



Photo by Sherrie Smith, University of Arkansas Cooperative Extension

## Cotton Bacterial Blight- *Xanthomonas campestris* pv. *malvacearum*



Photo by Sherrie Smith, University of Arkansas Cooperative Extension



## Cotton Bacterial Blight- *Xanthomonas campestris* pv. *malvacearum*




Photo by Sherrie Smith, University of Arkansas  
Cooperative Extension

## Cotton Bacterial Blight- *Xanthomonas campestris* pv. *malvacearum*



Photo by Sherrie Smith, University of Arkansas  
Cooperative Extension

## Grape

Downy Mildew of grape occurs in production areas where it is warm and wet during vegetative growth of the vine. The causal agent, *Plasmopara viticola*, can attack all green parts of the plant. Leaf lesions are yellowish and oily, or angular, yellow to reddish brown, and limited by the veins. A dense, white, cottony growth of sporulation occurs on the underside of the leaf. Infected shoots curl into a shepherd's crook, and become white with sporulation, eventually turning brown and dying. These same symptoms can occur on petioles, tendrils, and young inflorescences. Leaves drop prematurely, reducing sugar content in the fruit and decreasing winter hardiness of the buds. Young berries are also highly susceptible. They appear grayish and become covered with downy, felty sporulation. Infected berries do not ripen normally, but remain firm, eventually dropping from the vine. As with all downy mildews, good soil drainage is essential. The cleanup of fallen leaves and berries, and the removal of infected shoots helps limit inoculum, but rarely are these measures sufficient in themselves to control Downy Mildew in areas with high disease pressure. Fungicides must be applied, starting at 3-6" shoot growth. Captan, Mancozeb, Ziram, Abound, Sovran, Pristine, Aliette, Scala, Rovral, Reason, and Gavel are labeled for Downy Mildew control.



## **Grape Downy Mildew-*Plasmopara viticola***




Photo by Sherrie Smith, University of Arkansas Cooperative Extension

## **Grape Downy Mildew-*Plasmopara viticola***



Photo by Sherrie Smith, University of Arkansas Cooperative Extension

## **Grape Downy Mildew-*Plasmopara viticola***



Photo by Sherrie Smith, University of Arkansas Cooperative Extension

This bulletin from the Cooperative Extension Plant Health Clinic (Plant Disease Clinic) is an electronic update about diseases and other problems observed in our lab each month. Input from everybody interested in plants is welcome and appreciated.

"This work is supported by the Crop Protection and Pest Management Program [grant no. 2017-70006-27279/project accession no. 1013890] from the USDA National Institute of Food and Agriculture."

The University of Arkansas System Division of Agriculture offers all its Extension and Research programs to all eligible persons without regard to race, color, sex, gender identity, sexual orientation, national origin, religion, age, disability, marital or veteran status, genetic information, or any other legally protected status, and is an Affirmative Action/Equal Opportunity Employer.