



Sherrie Smith
Keiddy Urrea

Issue 27, August 15, 2017

Arkansas Plant Health Clinic Newsletter

Follow us on social media



Mulberry

Ornamental weeping mulberries are popular small trees, grown for their small size and interesting shape. Fruiting mulberries are grown around the world for their fruit, lumber, and for silkworm production. The Plant Health Clinic receives samples of mulberry leaves at this time of year with a fungal leaf spot caused by *Cercospora mori* or *Cercosporella mori*. The beginning symptoms of this disease are small dark spots in early spring that gradually increase in size through the growing season. The spots gradually become circular with the center appearing as a grayish white to tan spot with dark brown margins. Spores develop in the lesions during periods of wet weather and high humidity. Severe infections cause defoliation, which can weaken a tree already under stress. Weeping mulberries are small enough to be easily sprayed. Clean up all fallen leaves and spray with an ornamental fungicide. Homeowners may use Fertilome Broad Spectrum Lawn and Garden Fungicide, (chlorothalonil), or Hi-Yield Vegetable, Flower, Fruit, and Ornamental Fungicide, (chlorothalonil) or Ortho Maxx Garden Disease Control, (chlorothalonil), or Ortho Disease B Gon Garden Fungicide, (chlorothalonil), or

Garden Tech Daconil Fungicide, (chlorothalonil), or Bonide Fung-onil Multipurpose Fungicide, (chlorothalonil), or Spectracide Immunox Plus, (myclobutanil & permethrin). It is best to spray trees with a history of the disease early in the season before the spots develop.

Mulberry Leaf Spot-*Cercosporella mori*



Photo by Sherrie Smith, University of Arkansas Cooperative Extension

The University of Arkansas System Division of Agriculture offers all its Extension and Research programs to all eligible persons without regard to race, color, sex, gender identity, sexual orientation, national origin, religion, age, disability, marital or veteran status, genetic information, or any other legally protected status, and is an Affirmative Action/Equal Opportunity Employer.



Mulberry Leaf Spot-*Cercospora mori*



Photo by Brad McGinley, University of Arkansas Cooperative Extension

Lettuce

The collapse of lettuce plants caused by the oomycete *Pythium* can cause serious crop losses. Symptoms develop on lettuce that is at the rosette stage or older. Infected plants will be stunted and lag healthy lettuce. As the disease progresses, outer leaves will start to wilt during the warmer times of the day and eventually turn yellow before becoming brown and dead. In advanced stages, the entire plant can wilt and die. Below ground, the pathogen first attacks the small feeder roots of the lettuce, making them soft and brownish gray in color. Later in disease development, the taproot will also be darkly discolored, and the entire root system can be rotted. Secondary soft rot pathogens such as *Pectobacterium carotovorum*, may move into the damaged areas and exacerbate the rot.

The University of Arkansas System Division of Agriculture offers all its Extension and Research programs to all eligible persons without regard to race, color, sex, gender identity, sexual orientation, national origin, religion, age, disability, marital or veteran status, genetic information, or any other legally protected status, and is an Affirmative Action/Equal Opportunity Employer.

Actinovate may be used in organic systems and hydroponic systems. Ridomil Gold SL may be used in conventional systems. In conventional plantings, improve drainage and do not over water.

Lettuce Wilt-*Pythium* spp.



Photo by Sherrie Smith, University of Arkansas Cooperative Extension



Lettuce Wilt-*Pythium* spp.



Photo by Sherrie Smith, University of Arkansas Cooperative Extension

Lettuce Soft Rot-*Pectobacterium carotovorum* formerly *Erwinia carotovora*



Photo by Sherrie Smith, University of Arkansas Cooperative Extension

Canna

Two species of canna leaf roller are present in Arkansas, the Larger Canna Leaf Roller, *Calpodes ethlius* (Stoll) (Insecta: Lepidoptera: Hesperidae), and the Lesser Canna Leaf Roller, *Geshna cannalis* (Quaintance) (Insecta: Lepidoptera: Pyralidae). The Plant Health Clinic receives more samples of Lesser Canna Leaf Roller than Greater.

Canna leaf rollers, as their name implies, roll canna leaves, and keep them rolled using silk produced by larva (caterpillar). This is accomplished by attaching silk before the leaf unrolls (preferred by the lesser canna leaf roller) or by attaching silk to one edge of the leaf and pulling toward the other edge (usually the larger canna leaf roller). Once rolled, the leaf provides a protective area for the caterpillar to feed. The lesser canna leaf roller caterpillars generally feed on the surface of the leaf and do not chew completely through the leaf, but the larger leaf roller feeds through the leaf. Also, the Larger leafroller does not accumulate frass inside the rolled leaf but flicks it away. Later when the leaf opens, the feeding damage appears as holes in the leaves and ragged leaf edges.

Canna leaf rollers only feed on plants in the Genus *Canna* and close relatives, so other plants in the garden are not threatened. This also makes management a little easier because dead plant material can be cut to the ground in winter and disposed of which reduces the number of overwintering larvae and pupae. Systemic insecticides such as Merit or BioAdvanced Tree and Shrub Systemic Insecticide may be applied. Alternatively, a



Sherrie Smith
Keiddy Urrea

product applied to the leaves containing *Bacillus thuringiensis* (BT) gives control without toxicity to organisms other than members of the Lepidoptera.

Larger Canna Leaf Roller-*Calpodes ethlius*



Photo by Sherrie Smith, University of Arkansas Cooperative Extension

Lesser Canna Leaf Roller (note the silk ties)-*Geshna cannalis*



Photos by Sherrie Smith, University of Arkansas Cooperative Extension

The University of Arkansas System Division of Agriculture offers all its Extension and Research programs to all eligible persons without regard to race, color, sex, gender identity, sexual orientation, national origin, religion, age, disability, marital or veteran status, genetic information, or any other legally protected status, and is an Affirmative Action/Equal Opportunity Employer.



Sherrie Smith
Keiddy Urrea

This bulletin from the Cooperative Extension Plant Health Clinic (Plant Disease Clinic) is an electronic update about diseases and other problems observed in our lab each month. Input from everybody interested in plants is welcome and appreciated.

"This work is supported by the Crop Protection and Pest Management Program [grant no. 2017-70006-27279/project accession no. 1013890] from the USDA National Institute of Food and Agriculture."